

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 31/44, C07D 471/04</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/27929</b> <b>(43) International Publication Date:</b> 10 June 1999 (10.06.99)
<b>(21) International Application Number:</b> PCT/US98/25324 <b>(22) International Filing Date:</b> 30 November 1998 (30.11.98) <b>(30) Priority Data:</b> 08/984,732 4 December 1997 (04.12.97) US <b>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application</b> US 08/984,732 (CON) Filed on 4 December 1997 (04.12.97) <b>(71) Applicant (for all designated States except US):</b> MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> HALCZENKO, Wasyl [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). STUMP, Craig, A. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). <b>(74) Common Representative:</b> MERCK & CO., INC.; Daniel, Mark, R., 126 East Lincoln Avenue, Rahway, NJ 07065 (US).		<b>(81) Designated States:</b> AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> INHIBITORS OF FARNESYL-PROTEIN TRANSFERASE  <b>(57) Abstract</b>  The present invention is directed to compounds which inhibit farnesyl-protein transferase (FTase) and the farnesylation of the oncogene protein Ras. The invention is further directed to chemotherapeutic compositions containing the compounds of this invention and methods for inhibiting farnesyl-protein transferase and the farnesylation of the oncogene protein Ras.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE OF THE INVENTION

## INHIBITORS OF FARNESYL-PROTEIN TRANSFERASE

BACKGROUND OF THE INVENTION

5           The Ras protein is part of a signalling pathway that links cell surface growth factor receptors to nuclear signals initiating cellular proliferation. Biological and biochemical studies of Ras action indicate that Ras functions like a G-regulatory protein. In the inactive state, Ras is bound to GDP. Upon growth factor receptor activation Ras is  
10 induced to exchange GDP for GTP and undergoes a conformational change. The GTP-bound form of Ras propagates the growth stimulatory signal until the signal is terminated by the intrinsic GTPase activity of Ras, which returns the protein to its inactive GDP bound form (D.R. Lowy and D.M. Willumsen, *Ann. Rev. Biochem.* 62:851-  
15 891 (1993)). Mutated *ras* genes are found in many human cancers, including colorectal carcinoma, exocrine pancreatic carcinoma, and myeloid leukemias. The protein products of these genes are defective in their GTPase activity and constitutively transmit a growth stimulatory signal.

20           Ras must be localized to the plasma membrane for both normal and oncogenic functions. At least 3 post-translational modifications are involved with Ras membrane localization, and all 3 modifications occur at the C-terminus of Ras. The Ras C-terminus contains a sequence motif termed a "CAAX" or "Cys-Aaa<sup>1</sup>-Aaa<sup>2</sup>-Xaa" box (Cys is cysteine, Aaa is an aliphatic amino acid, the Xaa is any  
25 amino acid) (Willumsen *et al.*, *Nature* 310:583-586 (1984)). Depending on the specific sequence, this motif serves as a signal sequence for the enzymes farnesyl-protein transferase or geranylgeranyl-protein transferase, which catalyze the alkylation of the  
30 cysteine residue of the CAAX motif with a C<sub>15</sub> or C<sub>20</sub> isoprenoid, respectively. (S. Clarke., *Ann. Rev. Biochem.* 61:355-386 (1992); W.R. Schafer and J. Rine, *Ann. Rev. Genetics* 30:209-237 (1992)). The Ras protein is one of several proteins that are known to undergo post-translational farnesylation. Other farnesylated proteins include the Ras-

related GTP-binding proteins such as Rho, fungal mating factors, the nuclear lamins, and the gamma subunit of transducin. James, et al., *J. Biol. Chem.* 269, 14182 (1994) have identified a peroxisome associated protein Pxf which is also farnesylated. James, et al., have also suggested  
5 that there are farnesylated proteins of unknown structure and function in addition to those listed above.

Inhibition of farnesyl-protein transferase has been shown to block the growth of Ras-transformed cells in soft agar and to modify other aspects of their transformed phenotype. It has also been  
10 demonstrated that certain inhibitors of farnesyl-protein transferase selectively block the processing of the Ras oncoprotein intracellularly (N.E. Kohl *et al.*, *Science*, 260:1934-1937 (1993) and G.L. James *et al.*, *Science*, 260:1937-1942 (1993). Recently, it has been shown that an inhibitor of farnesyl-protein transferase blocks the growth of *ras*-  
15 dependent tumors in nude mice (N.E. Kohl *et al.*, *Proc. Natl. Acad. Sci U.S.A.*, 91:9141-9145 (1994) and induces regression of mammary and salivary carcinomas in *ras* transgenic mice (N.E. Kohl *et al.*, *Nature Medicine*, 1:792-797 (1995).

Indirect inhibition of farnesyl-protein transferase *in vivo*  
20 has been demonstrated with lovastatin (Merck & Co., Rahway, NJ) and compactin (Hancock *et al.*, *ibid*; Casey *et al.*, *ibid*; Schafer *et al.*, *Science* 245:379 (1989)). These drugs inhibit HMG-CoA reductase, the rate limiting enzyme for the production of polyisoprenoids including farnesyl pyrophosphate. Farnesyl-protein  
25 transferase utilizes farnesyl pyrophosphate to covalently modify the Cys thiol group of the Ras CAAX box with a farnesyl group (Reiss *et al.*, *Cell*, 62:81-88 (1990); Schaber *et al.*, *J. Biol. Chem.*, 265:14701-14704 (1990); Schafer *et al.*, *Science*, 249:1133-1139 (1990); Manne *et al.*, *Proc. Natl. Acad. Sci USA*, 87:7541-7545 (1990)). Inhibition of  
30 farnesyl pyrophosphate biosynthesis by inhibiting HMG-CoA reductase blocks Ras membrane localization in cultured cells. However, direct inhibition of farnesyl-protein transferase would be more specific and attended by fewer side effects than would occur with the required dose of a general inhibitor of isoprene biosynthesis.

Inhibitors of farnesyl-protein transferase (FPTase) have been described in two general classes. The first are analogs of farnesyl diphosphate (FPP), while the second class of inhibitors is related to the protein substrates (e.g., Ras) for the enzyme. The peptide derived inhibitors that have been described are generally cysteine containing molecules that are related to the CAAX motif that is the signal for protein prenylation. (Schaber *et al.*, *ibid*; Reiss *et al.*, *ibid*; Reiss *et al.*, *PNAS*, 88:732-736 (1991)). Such inhibitors may inhibit protein prenylation while serving as alternate substrates for the farnesyl-protein transferase enzyme, or may be purely competitive inhibitors (U.S. Patent 5,141,851, University of Texas; N.E. Kohl *et al.*, *Science*, 260:1934-1937 (1993); Graham, *et al.*, *J. Med. Chem.*, 37, 725 (1994)). In general, deletion of the thiol from a CAAX derivative has been shown to dramatically reduce the inhibitory potency of the compound. However, the thiol group potentially places limitations on the therapeutic application of FPTase inhibitors with respect to pharmacokinetics, pharmacodynamics and toxicity. Therefore, a functional replacement for the thiol is desirable.

It has recently been shown that farnesyl-protein transferase inhibitors are inhibitors of proliferation of vascular smooth muscle cells and are therefore useful in the prevention and therapy of arteriosclerosis and diabetic disturbance of blood vessels (JP H7-112930). It has also recently been disclosed that certain 1,2,3,4-tetrahydroisoquinoline peptidomimetic compounds, some of which incorporate an imidazole moiety, are inhibitors of FPTase (U.S. Pat. No. 5,439,918, EP 0 618 221 A2 and EP 0 675 112 A1 ).



It is, therefore, an object of this invention to develop novel peptidomimetic compounds that do not have a thiol moiety, and that will inhibit farnesyl-protein transferase and thus, the post-translational farnesylation of proteins. It is a further object of this invention to develop chemotherapeutic compositions containing the compounds of this invention and methods for producing the compounds of this invention.

## SUMMARY OF THE INVENTION

The present invention comprises substituted [4.3.0]-nitrogen containing ring systems and homologous compounds which inhibit farnesyl-protein transferase. Furthermore, these compounds differ from such heterocyclic compounds previously described as inhibitors of farnesyl-protein transferase with respect to the alkyl or heteroatom containing linker between the ring-nitrogen and the imidazolyl moiety, and with respect to the lack of a thiol moiety in the instant compounds. Further contained in this invention are

chemotherapeutic compositions containing these farnesyl transferase inhibitors and methods for their production.


The compounds of this invention are illustrated by the formulae I and A:



## DETAILED DESCRIPTION OF THE INVENTION

The compounds of this invention are useful in the inhibition of farnesyl-protein transferase and the farnesylation of the oncogene

protein Ras. In a first embodiment of this invention, the inhibitors of farnesyl-protein transferase are illustrated by the formula I:



5 wherein:

R¹ᵃ, R¹ᵇ and R¹ᶜ are independently selected from:

- a) hydrogen,
- b) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, R<sup>8</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,
- c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by unsubstituted or substituted aryl, heterocyclic, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, R<sup>8</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)-NR<sup>8</sup>-,

provided that R¹ᵃ is not unsubstituted or substituted imidazolyl;

- 20 R²ᵃ, R²ᵇ' and R²ᵇ'' are independently hydrogen or -(CR¹¹₂)<sub>v</sub>A³(CR¹²₂)<sub>w</sub>R¹³; or R²ᵇ' and R²ᵇ'' are combined as O;

R³ᵃ and R³ᵇ are independently selected from:

- 25 a) hydrogen,

- 5                   b)    unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, unsubstituted or substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl, unsubstituted or substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, halogen, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>9</sup>C(O)O-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, NO<sub>2</sub>, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,
- 10                  c)    unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl,
- d)    substituted C<sub>1</sub>-C<sub>6</sub> alkyl wherein the substituent on the substituted C<sub>1</sub>-C<sub>6</sub> alkyl is selected from unsubstituted or substituted aryl, unsubstituted or substituted heterocyclic, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, and
- 15                   R<sup>9</sup>OC(O)-NR<sup>8</sup>-;

R<sup>4</sup> is independently selected from:

- 20                  a)    hydrogen,
- b)    unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, perfluoroalkyl, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, R<sup>8</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-, and
- 25                  c)    C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl, heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, perfluoroalkyl, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NH-, CN, H<sub>2</sub>N-C(NH)-, R<sup>8</sup>C(O)-, R<sup>8</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>8</sup>OC(O)NH-,

provided that R<sup>4</sup> is not unsubstituted or substituted imidazolyl;

30

R<sup>5</sup> is independently selected from:

- a)    hydrogen,






- b) C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, perfluoroalkyl, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, (R<sup>8</sup>)<sub>2</sub>N-C-(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, R<sup>8</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-, and
- 5 c) C<sub>1</sub>-C<sub>6</sub> alkyl, unsubstituted or substituted by perfluoroalkyl, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, R<sup>8</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-;
- 10 R<sup>8</sup> is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, 2,2,2-trifluoroethyl and aryl;

R<sup>9</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

- 15 R<sup>10</sup> is selected from: H; R<sup>8</sup>C(O)-; R<sup>9</sup>S(O)<sub>m</sub>-; unsubstituted or substituted C<sub>1</sub>-4 alkyl, unsubstituted or substituted C<sub>3</sub>-6 cycloalkyl, unsubstituted or substituted heterocycle, unsubstituted or substituted aryl, substituted aroyl, unsubstituted or substituted heteroaroyl, substituted arylsulfonyl, unsubstituted or substituted heteroarylsulfonyl,
- 20 wherein the substituted group is substituted with one or two substituents selected from:

- a) C<sub>1</sub>-4 alkoxy,  
 b) aryl or heterocycle,  
 c) halogen,  
 25 d) HO,



- h)  $N(R^8)_2$ , or
- i) C3-6 cycloalkyl;

5 R<sup>11</sup> and R<sup>12</sup> are independently selected from:

- a) hydrogen,
- b) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by C<sub>2</sub>-C<sub>20</sub> alkenyl, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, N<sub>3</sub>, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,
- 10 c) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>20</sub> alkenyl, halogen, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-, and
- 15 d) C<sub>1</sub>-C<sub>6</sub> alkyl substituted with an unsubstituted or substituted group selected from aryl, heterocyclic and C<sub>3</sub>-C<sub>10</sub> cycloalkyl;

R<sup>13</sup> is selected from:

- 20 a) hydrogen,
- b) substituted or unsubstituted aryl, substituted or unsubstituted heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>2</sub>-C<sub>20</sub> alkynyl, C<sub>1</sub>-C<sub>20</sub> perfluoroalkyl, allyloxy, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, (R<sup>9</sup>)<sub>2</sub>NC(O)- or R<sup>9</sup>OC(O)NR<sup>8</sup>-, and
- 25 c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by substituted or unsubstituted aryl, substituted or unsubstituted heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>2</sub>-C<sub>20</sub> alkynyl, C<sub>2</sub>-C<sub>20</sub> perfluoroalkyl, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NH-, CN, H<sub>2</sub>N-C(NH)-, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NH-;
- 30

A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond, -CH=CH-, -C≡C-,  
 -C(O)-, -C(O)NR<sup>8</sup>-, -NR<sup>8</sup>C(O)-, O, -N(R<sup>8</sup>)-,  
 -S(O)<sub>2</sub>N(R<sup>8</sup>)-, -N(R<sup>8</sup>)S(O)<sub>2</sub>-, or -S(O)<sub>m</sub>;

5 A<sup>3</sup> are independently selected from: a bond, -CH=CH-,  
 -C≡C-, -C(O)-, -C(O)NR<sup>10</sup>-, -NR<sup>10</sup>C(O)-, O, -N(R<sup>10</sup>)-,  
 -S(O)<sub>2</sub>N(R<sup>10</sup>)-, -N(R<sup>10</sup>)S(O)<sub>2</sub>-, or -S(O)<sub>m</sub>;

J, K and L are independently selected from: S, N, O, NH and CH;

10

V is selected from:

- a) hydrogen,
- b) heterocycle,
- c) aryl,
- 15 d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are  
 replaced with a heteroatom selected from O, S, and N,  
 and
- e) C<sub>2</sub>-C<sub>20</sub> alkenyl,

provided that V is not hydrogen if A<sup>1</sup> is -S(O)<sub>m</sub> and V is not hydrogen

20 if A<sup>1</sup> is a bond, n is 0 and A<sup>2</sup> is -S(O)<sub>m</sub>;  
 and provided that V is not imidazolyl;

W is a heterocycle;

25 X is a bond, -S(O)<sub>m</sub>-, O or -C(=O)-;

m is 0, 1 or 2;  
 n is 0, 1, 2, 3 or 4;  
 p is 0, 1, 2, 3 or 4;  
 30 q is 0, 1, 2, 3 or 4, provided that q is not 0 or 1 if X is O;  
 r is 0 to 5, provided that r is 0 when V is hydrogen;  
 s is 1 or 2;  
 t is 0 or 1;  
 u is independently 0, 1 or 2;

v is 0, 1, 2, 3 or 4, provided that v is not 0 when A<sup>3</sup> is  
 -NR<sup>10</sup>C(O)-, O, -N(R<sup>10</sup>)-, -S(O)<sub>2</sub>N(R<sup>10</sup>)-,  
 -N(R<sup>10</sup>)S(O)<sub>2</sub>-, or S(O)<sub>m</sub>;


w is 0, 1, 2, 3 or 4; and

5

the dashed lines represent optional double bonds;

or an optical isomer or a pharmaceutically acceptable salt thereof.

10 In another embodiment of this invention, the inhibitors  
 of farnesyl-protein transferase are illustrated by the formula A:



wherein:

15 R<sup>1a</sup>, R<sup>1b</sup> and R<sup>1c</sup> are independently selected from:

- a) hydrogen,
- b) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, R<sup>8</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,
- 20 c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by unsubstituted or substituted aryl, heterocyclic, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, R<sup>8</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)-NR<sup>8</sup>-,
- 25

provided that R<sup>1a</sup> is not unsubstituted or substituted imidazolyl;

R<sup>2a</sup>, R<sup>2b'</sup> and R<sup>2b''</sup> are independently hydrogen or  
 -(CR<sup>11</sup><sub>2</sub>)<sub>v</sub>A<sup>3</sup>(CR<sup>12</sup><sub>2</sub>)<sub>w</sub>R<sup>13</sup>; or

5 R<sup>2b'</sup> and R<sup>2b''</sup> are combined as O;

R<sup>3a</sup> and R<sup>3b</sup> are independently selected from:

- a) hydrogen,
- 10 b) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, unsubstituted or substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl, unsubstituted or substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, halogen, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>9</sup>C(O)O-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, NO<sub>2</sub>, R<sup>8</sup>C(O)-, N<sub>3</sub>,  
 15 -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,
- c) unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl,
- 20 d) substituted C<sub>1</sub>-C<sub>6</sub> alkyl wherein the substituent on the substituted C<sub>1</sub>-C<sub>6</sub> alkyl is selected from unsubstituted or substituted aryl, unsubstituted or substituted heterocyclic, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, and R<sup>9</sup>OC(O)-NR<sup>8</sup>-;

25 R<sup>4</sup> is independently selected from:

- a) hydrogen,
- 30 b) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, perfluoroalkyl, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, R<sup>8</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-, and
- c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl, heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub>

alkynyl, perfluoroalkyl, F, Cl, Br,  $R^8O-$ ,  $R^9S(O)_m-$ ,  
 $R^8C(O)NH-$ , CN,  $H_2N-C(NH)-$ ,  $R^8C(O)-$ ,  $R^8OC(O)-$ ,  $N_3$ ,  
 $-N(R^8)_2$ , or  $R^8OC(O)NH-$ ,

provided that  $R^4$  is not unsubstituted or substituted imidazolyl;

5

$R^5$  is independently selected from:




- a) hydrogen,
- b)  $C_2-C_6$  alkenyl,  $C_2-C_6$  alkynyl,  $C_3-C_6$  cycloalkyl,  
 perfluoroalkyl, F, Cl, Br,  $R^8O-$ ,  $R^9S(O)_m-$ ,  $R^8C(O)NR^8-$ ,  
 10 CN,  $NO_2$ ,  $(R^8)_2N-C-(NR^8)-$ ,  $R^8C(O)-$ ,  $R^8OC(O)-$ ,  $N_3$ ,  
 $-N(R^8)_2$ , or  $R^9OC(O)NR^8-$ , and
- c)  $C_1-C_6$  alkyl, unsubstituted or substituted by perfluoroalkyl,  
 F, Cl, Br,  $R^8O-$ ,  $R^9S(O)_m-$ ,  $R^8C(O)NR^8-$ , CN,  $(R^8)_2N-$   
 $C(NR^8)-$ ,  $R^8C(O)-$ ,  $R^8OC(O)-$ ,  $N_3$ ,  $-N(R^8)_2$ , or  
 15  $R^9OC(O)NR^8-$ ;

$R^8$  is independently selected from hydrogen,  $C_1-C_6$  alkyl, benzyl, 2,2,2-trifluoroethyl and aryl;

20  $R^9$  is independently selected from  $C_1-C_6$  alkyl and aryl;

$R^{10}$  is selected from: H;  $R^8C(O)-$ ;  $R^9S(O)_m-$ ; unsubstituted or substituted  $C_1-4$  alkyl, unsubstituted or substituted  $C_3-6$  cycloalkyl, unsubstituted or substituted heterocycle, unsubstituted or substituted  
 25 aryl, substituted aroyl, unsubstituted or substituted heteroaroyl, substituted arylsulfonyl, unsubstituted or substituted heteroarylsulfonyl, wherein the substituted group is substituted with one or two substituents selected from:

- a)  $C_1-4$  alkoxy,
- 30 b) aryl or heterocycle,
- c) halogen,
- d) HO,



h)  $\text{N}(\text{R}^8)_2$ , or

i) C3-6 cycloalkyl;

5

R11 and R12 are independently selected from:

- a) hydrogen,
- b) C1-C6 alkyl unsubstituted or substituted by C2-C20 alkenyl,  $\text{R}^8\text{O}-$ ,  $\text{R}^9\text{S}(\text{O})_m-$ ,  $\text{R}^8\text{C}(\text{O})\text{NR}^8-$ , CN, N3, (R8)2N-C(NR8)-,  $\text{R}^8\text{C}(\text{O})-$ ,  $-\text{N}(\text{R}^8)_2$ , or  $\text{R}^9\text{OC}(\text{O})\text{NR}^8-$ ,
- 10 c) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, C3-C10 cycloalkyl, C2-C20 alkenyl, halogen,  $\text{R}^8\text{O}-$ ,  $\text{R}^9\text{S}(\text{O})_m-$ ,  $\text{R}^8\text{C}(\text{O})\text{NR}^8-$ , CN, NO2, (R8)2N-C(NR8)-,  $\text{R}^8\text{C}(\text{O})-$ , N3,  $-\text{N}(\text{R}^8)_2$ , or  $\text{R}^9\text{OC}(\text{O})\text{NR}^8-$ , and
- 15 d) C1-C6 alkyl substituted with an unsubstituted or substituted group selected from aryl, heterocyclic and C3-C10 cycloalkyl;

R<sup>13</sup> is selected from:

- a) hydrogen,
- b) substituted or unsubstituted aryl, substituted or unsubstituted heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>2</sub>-C<sub>20</sub> alkynyl, C<sub>1</sub>-C<sub>20</sub> perfluoroalkyl, allyloxy, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, (R<sup>9</sup>)<sub>2</sub>NC(O)- or R<sup>9</sup>OC(O)NR<sup>8</sup>-, and
- c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by substituted or unsubstituted aryl, substituted or unsubstituted heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>2</sub>-C<sub>20</sub> alkynyl, C<sub>2</sub>-C<sub>20</sub> perfluoroalkyl, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NH-, CN, H<sub>2</sub>N-C(NH)-, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NH-;

15

A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR<sup>8</sup>-, -NR<sup>8</sup>C(O)-, O, -N(R<sup>8</sup>)-, -S(O)<sub>2</sub>N(R<sup>8</sup>)-, -N(R<sup>8</sup>)S(O)<sub>2</sub>-, or -S(O)<sub>m</sub>;

- 20 A<sup>3</sup> are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR<sup>10</sup>-, -NR<sup>10</sup>C(O)-, O, -N(R<sup>10</sup>)-, -S(O)<sub>2</sub>N(R<sup>10</sup>)-, -N(R<sup>10</sup>)S(O)<sub>2</sub>-, or -S(O)<sub>m</sub>;

J, K and L are independently selected from: S, N, O, NH and CH;

25

V is selected from:

- a) hydrogen,
- b) heterocycle,
- c) aryl,
- d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
- e) C<sub>2</sub>-C<sub>20</sub> alkenyl,



provided that V is not hydrogen if  $A^1$  is  $-S(O)_m$  and V is not hydrogen if  $A^1$  is a bond, n is 0 and  $A^2$  is  $-S(O)_m$ ; and provided that V is not imidazolyl;

5 W is a heterocycle;

X is a bond,  $-S(O)_m-$ , O or  $-C(=O)-$ ;


- m is 0, 1 or 2;
- 10 n is 0, 1, 2, 3 or 4;
- p is 0, 1, 2, 3 or 4;
- q is 0, 1, 2, 3 or 4, provided that q is not 0 or 1 if X is O;
- r is 0 to 5, provided that r is 0 when V is hydrogen;
- s is 1 or 2;
- 15 t is 0 or 1;
- u is independently 0, 1 or 2;
- v is 0, 1, 2, 3 or 4, provided that v is not 0 when  $A^3$  is  $-NR^{10}C(O)-$ , O,  $-N(R^{10})-$ ,  $-S(O)_2N(R^{10})-$ ,  $-N(R^{10})S(O)_2-$ , or  $S(O)_m$ ;
- 20 w is 0, 1, 2, 3 or 4; and

the dashed lines represent optional double bonds;

or an optical isomer or a pharmaceutically acceptable salt thereof.

25

A preferred embodiment of the compounds of this invention is illustrated by the following formula:




wherein:

R<sup>1a</sup> and R<sup>1c</sup> are independently selected from: hydrogen, C<sub>3</sub>-C<sub>10</sub>  
 5 cycloalkyl, R<sup>8</sup>O-, -N(R<sup>8</sup>)<sub>2</sub>, F or C<sub>1</sub>-C<sub>6</sub> alkyl;

R<sup>1b</sup> is independently selected from:

- a) hydrogen,
- b) unsubstituted or substituted aryl, unsubstituted or  
 10 substituted heterocycle, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, R<sup>8</sup>O-, -N(R<sup>8</sup>)<sub>2</sub>  
 or C<sub>2</sub>-C<sub>6</sub> alkenyl,
- c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by unsubstituted or  
 substituted aryl, heterocycle, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub>  
 alkenyl, R<sup>8</sup>O-, or -N(R<sup>8</sup>)<sub>2</sub>;

15 R<sup>2a</sup> and R<sup>2b'</sup> are independently selected from selected from:  
 H; C<sub>1</sub>-C<sub>6</sub> alkyl,



20 R<sup>3a</sup> and R<sup>3b</sup> are independently selected from:

- a) hydrogen,
- b) unsubstituted or substituted aryl, unsubstituted or  
 substituted heterocycle, unsubstituted or substituted  
 C<sub>3</sub>-C<sub>10</sub> cycloalkyl, unsubstituted or substituted C<sub>2</sub>-C<sub>6</sub>  
 25 alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, halogen, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,

- $R^9O-$ ,  $R^9S(O)_m-$ ,  $R^8C(O)NR^8-$ ,  $(R^8)_2NC(O)-$ ,  $R^9C(O)O-$ ,  
 $R^8_2N-C(NR^8)-$ ,  $CN$ ,  $NO_2$ ,  $R^8C(O)-$ ,  $N_3$ ,  $-N(R^8)_2$ , or  
 $R^9OC(O)NR^8-$ ,  
 5 c) unsubstituted C1-C6 alkyl,  
 d) substituted C1-C6 alkyl wherein the substituent on the  
 substituted C1-C6 alkyl is selected from unsubstituted or  
 substituted aryl, unsubstituted or substituted heterocyclic,  
 C3-C10 cycloalkyl, C2-C6 alkenyl, C2-C6 alkynyl,  $R^9O-$ ,  
 $R^9S(O)_m-$ ,  $R^8C(O)NR^8-$ ,  $(R^8)_2NC(O)-$ ,  $R^8_2N-C(NR^8)-$ ,  
 10  $CN$ ,  $R^8C(O)-$ ,  $N_3$ ,  $-N(R^8)_2$ , and  $R^9OC(O)-NR^8-$ ;

$R^4$  is independently selected from:

- a) hydrogen,  
 b) C1-C6 alkyl, C2-C6 alkenyl, C2-C6 alkynyl, C1-C6  
 15 perfluoroalkyl,  $F$ ,  $Cl$ ,  $R^8O-$ ,  $R^8C(O)NR^8-$ ,  $CN$ ,  $NO_2$ ,  
 $(R^8)_2N-C(NR^8)-$ ,  $R^8C(O)-$ ,  $R^8OC(O)-$ ,  $-N(R^8)_2$ , or  
 $R^9OC(O)NR^8-$ , and  
 c) C1-C6 alkyl substituted by C1-C6 perfluoroalkyl,  $R^8O-$ ,  
 $R^8C(O)NR^8-$ ,  $(R^8)_2N-C(NR^8)-$ ,  $R^8C(O)-$ ,  $R^8OC(O)-$ ,  
 20  $-N(R^8)_2$ , or  $R^9OC(O)NR^8-$ ;

$R^5$  is selected from:

- a) hydrogen,  
 b) C2-C6 alkenyl, C2-C6 alkynyl, C3-C6 cycloalkyl, C1-C6  
 25 perfluoroalkyl,  $F$ ,  $Cl$ ,  $R^8O-$ ,  $R^9S(O)_m-$ ,  $R^8C(O)NR^8-$ ,  $CN$ ,  
 $NO_2$ ,  $(R^8)_2N-C(NR^8)-$ ,  $R^8C(O)-$ ,  $R^8OC(O)-$ ,  $-N(R^8)_2$ , or  
 $R^9OC(O)NR^8-$ , and  
 c) C1-C6 alkyl unsubstituted or substituted by C1-C6  
 perfluoroalkyl,  $F$ ,  $Cl$ ,  $R^8O-$ ,  $R^9S(O)_m-$ ,  $R^8C(O)NR^8-$ ,  $CN$ ,  
 30  $(R^8)_2N-C(NR^8)-$ ,  $R^8C(O)-$ ,  $R^8OC(O)-$ ,  $-N(R^8)_2$ , or  
 $R^9OC(O)NR^8-$ ;

R<sup>6</sup> and R<sup>7</sup> are independently selected from:

H; C<sub>1</sub>-4 alkyl, C<sub>3</sub>-6 cycloalkyl, aryl, heterocycle,  
unsubstituted or substituted with one or two:



- a) C<sub>1</sub>-4 alkoxy,
- b) halogen, or
- c) substituted or unsubstituted aryl or substituted  
or unsubstituted heterocycle,

R<sup>8</sup> is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, 2,2,2-trifluoroethyl and aryl;

R<sup>9</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

R<sup>10</sup> is selected from: H; R<sup>8</sup>C(O)-; R<sup>9</sup>S(O)<sub>m</sub>-; unsubstituted or substituted C<sub>1</sub>-4 alkyl, unsubstituted or substituted C<sub>3</sub>-6 cycloalkyl, unsubstituted or substituted heterocycle, unsubstituted or substituted aryl, substituted aroyl, unsubstituted or substituted heteroaroyl, substituted arylsulfonyl, unsubstituted or substituted heteroarylsulfonyl, wherein the substituted group is substituted with one or two substituents selected from:

- a) C<sub>1</sub>-4 alkoxy,
- b) aryl or heterocycle,
- c) halogen,
- d) HO,



g) —S(O)<sub>m</sub>R<sup>9</sup>,

h) N(R<sup>8</sup>)<sub>2</sub>, or

i) C<sub>3</sub>-6 cycloalkyl;

A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond, -CH=CH-, -C≡C-,  
-C(O)-, -C(O)NR<sup>8</sup>-, O, -N(R<sup>8</sup>)-, or -S(O)<sub>m</sub>;

J, K and L are independently selected from: S, N, O, NH and CH;

5

V is selected from:

- a) heterocycle selected from pyrrolidinyl, imidazolyl,  
pyridinyl, thiazolyl, indolyl, quinolinyl, isoquinolinyl, and  
thienyl, and  
10 b) aryl;

W is a heterocycle selected from pyrrolidinyl, triazolyl, imidazolyl,  
pyridinyl, thiazolyl, indolyl, quinolinyl, or isoquinolinyl;

15 X is a bond, -S(O)<sub>m</sub>-, O or -C(=O)-;

m is 0, 1 or 2;

n is 0, 1, 2, 3 or 4;

p is 1, 2 or 3;

20 q is 0, 1 or 2, provided that q is not 0 or 1 if X is O;

r is 0 to 5, provided that r is 0 when V is hydrogen;

s is 1 or 2;


t is 1; and

u is independently 0 or 1;

25

or an optical isomer or a pharmaceutically acceptable salt thereof.

Another preferred embodiment of the compounds of this invention are illustrated by the formula B:



wherein:

5

R<sup>1a</sup> and R<sup>1c</sup> are independently selected from: hydrogen, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, R<sup>8</sup>O-, -N(R<sup>8</sup>)<sub>2</sub>, F or C<sub>1</sub>-C<sub>6</sub> alkyl;


R<sup>1b</sup> is independently selected from:

10

- a) hydrogen,
- b) aryl, heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, R<sup>8</sup>O-, -N(R<sup>8</sup>)<sub>2</sub>, F or C<sub>2</sub>-C<sub>6</sub> alkenyl,
- c) unsubstituted or substituted C<sub>1</sub>-C<sub>6</sub> alkyl wherein the substituent on the substituted C<sub>1</sub>-C<sub>6</sub> alkyl is selected from unsubstituted or substituted aryl, heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, R<sup>8</sup>O- and -N(R<sup>8</sup>)<sub>2</sub>;

15

R<sup>2a</sup> and R<sup>2b'</sup> are independently selected from selected from: H; C<sub>1</sub>-C<sub>6</sub> alkyl,



20

R<sup>3a</sup> and R<sup>3b</sup> are independently selected from:

- a) hydrogen,

- 5                   b)    unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, unsubstituted or substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl, unsubstituted or substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, halogen, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>9</sup>C(O)O-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, NO<sub>2</sub>, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,
- 10                   c)    unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl,
- d)    substituted C<sub>1</sub>-C<sub>6</sub> alkyl wherein the substituent on the substituted C<sub>1</sub>-C<sub>6</sub> alkyl is selected from unsubstituted or substituted aryl, unsubstituted or substituted heterocyclic, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, and R<sup>9</sup>OC(O)-NR<sup>8</sup>-;
- 15                   R<sup>4</sup> is independently selected from:
- a)    hydrogen,
- b)    aryl, substituted aryl, heterocycle, substituted heterocycle, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>8</sup>O-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-, and
- 20                   c)    C<sub>1</sub>-C<sub>6</sub> alkyl substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>8</sup>O-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-;
- 25                   R<sup>5a</sup> and R<sup>5b</sup> are independently hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, cyclopropyl, trifluoromethyl and halogen;
- 30                   R<sup>6</sup> and R<sup>7</sup> are independently selected from:
- H; C<sub>1</sub>-4 alkyl, C<sub>3</sub>-6 cycloalkyl, aryl, heterocycle, unsubstituted or substituted with one or two:
- a)    C<sub>1</sub>-4 alkoxy,

- b) halogen, or
- c) substituted or unsubstituted aryl or substituted or unsubstituted heterocycle;

5  $R^8$  is independently selected from hydrogen,  $C_1$ - $C_6$  alkyl, 2,2,2-trifluoroethyl, benzyl and aryl;

$R^9$  is independently selected from  $C_1$ - $C_6$  alkyl and aryl;

10  $A^1$  and  $A^2$  are independently selected from: a bond,  $-CH=CH-$ ,  $-C\equiv C-$ ,  $-C(O)-$ ,  $-C(O)NR^8-$ , O,  $-N(R^8)-$ , or  $-S(O)_m$ ;

J, K and L are independently selected from: S, N, O, NH and CH;

15 V is selected from:

- a) hydrogen,
- b) heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, oxazolyl, indolyl, quinolinyl, isoquinolinyl, triazolyl and thienyl,
- 20 c) aryl,
- d)  $C_1$ - $C_{20}$  alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
- e)  $C_2$ - $C_{20}$  alkenyl, and

25 provided that V is not hydrogen if  $A^1$  is  $-S(O)_m$  and V is not hydrogen if  $A^1$  is a bond, n is 0 and  $A^2$  is  $-S(O)_m$ ;

X is a bond,  $-S(O)_m-$ , O or  $-C(=O)-$ ;


30 m is 0, 1 or 2;  
 n is 0, 1, 2, 3 or 4;  
 p is 0, 1, 2, 3 or 4;  
 q is 0, 1 or 2, provided that q is not 0 or 1 if X is O;  
 r is 0 to 5, provided that r is 0 when V is hydrogen; and



u is independently 0 or 1;

or an optical isomer or pharmaceutically acceptable salt thereof.


- 5 Another preferred embodiment of the compounds of this invention are illustrated by the formula C:



wherein:

- 10 R<sup>1a</sup> and R<sup>1c</sup> are independently selected from: hydrogen, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, R<sup>8</sup>O-, -N(R<sup>8</sup>)<sub>2</sub>, F or C<sub>1</sub>-C<sub>6</sub> alkyl;
- R<sup>1b</sup> is independently selected from:
- 15 a) hydrogen,
- b) aryl, heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, R<sup>8</sup>O-, -N(R<sup>8</sup>)<sub>2</sub>, F or C<sub>2</sub>-C<sub>6</sub> alkenyl,
- c) unsubstituted or substituted C<sub>1</sub>-C<sub>6</sub> alkyl wherein the substituent on the substituted C<sub>1</sub>-C<sub>6</sub> alkyl is selected from unsubstituted or substituted aryl, heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, R<sup>8</sup>O- and -N(R<sup>8</sup>)<sub>2</sub>;
- 20

R<sup>2a</sup> and R<sup>2b'</sup> are independently selected from selected from:  
H; C<sub>1</sub>-C<sub>6</sub> alkyl,



R<sup>3a</sup> and R<sup>3b</sup> are independently selected from:

- a) hydrogen,
- 5 b) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, unsubstituted or substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl, unsubstituted or substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, halogen, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>9</sup>C(O)O-,  
 10 R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, NO<sub>2</sub>, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,
- c) unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl,
- d) substituted C<sub>1</sub>-C<sub>6</sub> alkyl wherein the substituent on the substituted C<sub>1</sub>-C<sub>6</sub> alkyl is selected from unsubstituted or  
 15 substituted aryl, unsubstituted or substituted heterocyclic, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, and R<sup>9</sup>OC(O)-NR<sup>8</sup>-;

20 R<sup>4</sup> is independently selected from:

- a) hydrogen,
- b) aryl, substituted aryl, heterocycle, substituted heterocycle, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>8</sup>O-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>,  
 25 (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,  
 and
- c) C<sub>1</sub>-C<sub>6</sub> alkyl substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>8</sup>O-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-;

30

R<sup>5a</sup> and R<sup>5b</sup> are independently hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, cyclopropyl, trifluoromethyl and halogen;


- $R^6$  and  $R^7$  are independently selected from:  
H; C<sub>1</sub>-4 alkyl, C<sub>3</sub>-6 cycloalkyl, aryl, heterocycle,  
unsubstituted or substituted with one or two:
- 5      a) C<sub>1</sub>-4 alkoxy,  
      b) halogen, or  
      c) substituted or unsubstituted aryl or substituted  
          or unsubstituted heterocycle;
- 10     $R^8$  is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, 2,2,2-trifluoroethyl, benzyl and aryl;
- $R^9$  is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;
- 15     $A^1$  and  $A^2$  are independently selected from: a bond, -CH=CH-, -C≡C-,  
          -C(O)-, -C(O)NR<sup>8</sup>-, O, -N(R<sup>8</sup>)-, or -S(O)<sub>m</sub>;
- J, K and L are independently selected from: S, N, O, NH and CH;
- 20    V is selected from:
- a) hydrogen,  
      b) heterocycle selected from pyrrolidinyl, imidazolynyl,  
          pyridinyl, thiazolyl, oxazolyl, indolyl, quinolynyl,  
          isoquinolynyl, triazolyl and thienyl,  
25    c) aryl,  
      d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are  
          replaced with a heteroatom selected from O, S, and N,  
          and  
      e) C<sub>2</sub>-C<sub>20</sub> alkenyl, and
- 30    provided that V is not hydrogen if  $A^1$  is -S(O)<sub>m</sub> and V is not hydrogen  
      if  $A^1$  is a bond, n is 0 and  $A^2$  is -S(O)<sub>m</sub>;
- X is a bond, -S(O)<sub>m</sub>-, O or -C(=O)-;

- m is 0, 1 or 2;  
 n is 0, 1, 2, 3 or 4;  
 p is 0, 1, 2, 3 or 4, provided that p is not 0 if X is a bond or O;  
 5 q is 0, 1 or 2, provided that q is not 0 or 1 if X is O;  
 r is 0 to 5, provided that r is 0 when V is hydrogen; and  
 u is independently 0 or 1;

or an optical isomer or pharmaceutically acceptable salt thereof.

10

In a more preferred embodiment of this invention, the inhibitors of farnesyl-protein transferase are illustrated by the formula D:



15 wherein:

R1a and R1c are independently selected from: hydrogen, C3-C10 cycloalkyl or C1-C6 alkyl;


20 R1b is independently selected from:

- a) hydrogen,
  - b) aryl, heterocycle, C3-C10 cycloalkyl, R<sup>8</sup>O-, -N(R<sup>8</sup>)<sub>2</sub>, F or C<sub>2</sub>-C<sub>6</sub> alkenyl,
  - c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl,
- 25 heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, R<sup>8</sup>O-, or



R<sup>2a</sup> and R<sup>2b'</sup> are independently selected from selected from:

5 H; C<sub>1</sub>-C<sub>6</sub> alkyl,



R<sup>3a</sup> and R<sup>3b</sup> are independently selected from:

- 10 a) hydrogen,  
b) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, unsubstituted or substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl, unsubstituted or substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, halogen, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>9</sup>C(O)O-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-,  
15 CN, NO<sub>2</sub>, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,  
c) unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl,  
d) substituted C<sub>1</sub>-C<sub>6</sub> alkyl wherein the substituent on the substituted C<sub>1</sub>-C<sub>6</sub> alkyl is selected from unsubstituted or substituted aryl, unsubstituted or substituted heterocyclic, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, and R<sup>9</sup>OC(O)-NR<sup>8</sup>-;
- 20

25 R<sup>4</sup> is independently selected from:

- a) hydrogen,  
b) aryl, substituted aryl, heterocycle, substituted heterocycle, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>8</sup>O-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,  
30 and

- c) C<sub>1</sub>-C<sub>6</sub> alkyl substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>8</sup>O-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-;

5 R<sup>5a</sup> and R<sup>5b</sup> are independently hydrogen, ethyl, cyclopropyl or methyl;

R<sup>6</sup> and R<sup>7</sup> are independently selected from:

H; C<sub>1</sub>-4 alkyl, C<sub>3</sub>-6 cycloalkyl, aryl, heterocycle, unsubstituted or substituted with one or two:

- 10 a) C<sub>1</sub>-4 alkoxy,  
b) halogen, or  
c) substituted or unsubstituted aryl or substituted or unsubstituted heterocycle;

15 R<sup>8</sup> is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, 2,2,2-trifluoroethyl, benzyl and aryl;

R<sup>9</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

20 A<sup>1</sup> is selected from: a bond, -C(O)-, O, -N(R<sup>8</sup>)-, or -S(O)<sub>m</sub>;

J, K and L are independently selected from: S, N, O, NH and CH;

X is a bond, -S(O)<sub>m</sub>-, O or -C(=O)-;

25

n is 0, 1 or 2; provided that n is not 0 or 1 if A<sup>1</sup> is a bond, O, -N(R<sup>8</sup>)-, or S(O)<sub>m</sub>;

m is 0, 1 or 2;

p is 0, 1, 2, 3 or 4;


30 q is 0, 1 or 2, provided that q is not 0 or 1 if X is O;

r is 1 or 2; and

u is independently 0 or 1;

or an optical isomer or pharmaceutically acceptable salt thereof.

In another more preferred embodiment of this invention, the inhibitors of farnesyl-protein transferase are illustrated by the formula E:



5

wherein:

R<sup>1a</sup> and R<sup>1c</sup> are independently selected from: hydrogen, R<sup>8</sup>O-,  
-N(R<sup>8</sup>)<sub>2</sub>, F, C<sub>3</sub>-C<sub>10</sub> cycloalkyl or C<sub>1</sub>-C<sub>6</sub> alkyl;


10

R<sup>1b</sup> is independently selected from:

- a) hydrogen,
- b) aryl, heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, R<sup>8</sup>O-, -N(R<sup>8</sup>)<sub>2</sub>, F or C<sub>2</sub>-C<sub>6</sub> alkenyl,
- c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl,  
heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, R<sup>8</sup>O-, or  
-N(R<sup>8</sup>)<sub>2</sub>;

15

R<sup>2a</sup> and R<sup>2b'</sup> are independently selected from selected from:

20 H; C<sub>1</sub>-C<sub>6</sub> alkyl,

R<sup>3a</sup> and R<sup>3b</sup> are independently selected from:

- a) hydrogen,

- 5                   b)    unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, unsubstituted or substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl, unsubstituted or substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, halogen, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>9</sup>C(O)O-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, NO<sub>2</sub>, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,
- 10                   c)    unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl,
- d)    substituted C<sub>1</sub>-C<sub>6</sub> alkyl wherein the substituent on the substituted C<sub>1</sub>-C<sub>6</sub> alkyl is selected from unsubstituted or substituted aryl, unsubstituted or substituted heterocyclic, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, and R<sup>9</sup>OC(O)-NR<sup>8</sup>-;
- 15                   R<sup>4</sup> is independently selected from:
- a)    hydrogen,
- b)    aryl, substituted aryl, heterocycle, substituted heterocycle, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>8</sup>O-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-, and
- 20                   c)    C<sub>1</sub>-C<sub>6</sub> alkyl substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>8</sup>O-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-;
- 25

R<sup>5a</sup> and R<sup>5b</sup> are independently hydrogen, ethyl, cyclopropyl or methyl;

- 30                   R<sup>6</sup> and R<sup>7</sup> are independently selected from:  
                   H; C<sub>1</sub>-4 alkyl, C<sub>3</sub>-6 cycloalkyl, aryl, heterocycle, unsubstituted or substituted with one or two:
- a)    C<sub>1</sub>-4 alkoxy,
- b)    halogen, or



c) substituted or unsubstituted aryl or substituted or unsubstituted heterocycle;

5  $R^8$  is independently selected from hydrogen,  $C_1$ - $C_6$  alkyl, 2,2,2-trifluoroethyl, benzyl and aryl;

$R^9$  is independently selected from  $C_1$ - $C_6$  alkyl and aryl;

10 J, K and L are independently selected from: S, N, O, NH and CH;

X is a bond,  $-S(O)_m-$ , O or  $-C(=O)-$ ;

n is 0 or 1;

m is 0, 1 or 2;

15 p is 0, 1, 2, 3 or 4, provided that p is not 0 if X is a bond or O;


q is 0, 1 or 2, provided that q is not 0 or 1 if X is O;

r is 1 or 2; and

u is independently 0 or 1;

20 or an optical isomer or pharmaceutically acceptable salt thereof.

In a further embodiment of this invention, the inhibitors of farnesyl-protein transferase are illustrated by the formula F:



25

wherein:


R<sup>1a</sup> and R<sup>1c</sup> are independently selected from: hydrogen, C<sub>3</sub>-C<sub>10</sub> cycloalkyl or C<sub>1</sub>-C<sub>6</sub> alkyl;

5

R<sup>1b</sup> is independently selected from:

- a) hydrogen,
  - b) aryl, heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, R<sup>8</sup>O-, -N(R<sup>8</sup>)<sub>2</sub> or F,
  - c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl,
- 10 heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, R<sup>8</sup>O-, or -N(R<sup>8</sup>)<sub>2</sub>;

R<sup>2a</sup> and R<sup>2b'</sup> are independently selected from selected from: H; C<sub>1</sub>-C<sub>6</sub> alkyl,



15

R<sup>3a</sup> and R<sup>3b</sup> are independently selected from:

- a) hydrogen,
- b) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, unsubstituted or substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl, unsubstituted or substituted C<sub>2</sub>-C<sub>6</sub> alkenyl,
- 20 C<sub>2</sub>-C<sub>6</sub> alkynyl, halogen, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>9</sup>C(O)O-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, NO<sub>2</sub>, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,
- 25 c) unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl,
- d) substituted C<sub>1</sub>-C<sub>6</sub> alkyl wherein the substituent on the substituted C<sub>1</sub>-C<sub>6</sub> alkyl is selected from unsubstituted or substituted aryl, unsubstituted or substituted heterocyclic,
- 30 C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, and R<sup>9</sup>OC(O)-NR<sup>8</sup>-;

R<sup>5a</sup> and R<sup>5b</sup> are independently hydrogen, ethyl, cyclopropyl or methyl;

R<sup>6</sup> and R<sup>7</sup> are independently selected from:

- 5 H; C<sub>1</sub>-4 alkyl, C<sub>3</sub>-6 cycloalkyl, aryl, heterocycle,  
unsubstituted or substituted with one or two:  
a) C<sub>1</sub>-4 alkoxy,  
b) halogen, or  
c) substituted or unsubstituted aryl or substituted or  
unsubstituted heterocycle;

10

R<sup>8</sup> is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, 2,2,2-trifluoroethyl, benzyl and aryl;

R<sup>9</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

15

J, K and L are independently selected from: S, N, O, NH and CH;

X is a bond, -S(O)<sub>m</sub>-, O or -C(=O)-;

20

m is 0, 1 or 2;

p is 0, 1, 2, 3 or 4;


q is 0, 1 or 2, provided that q is not 0 or 1 if X is O; and

u is independently 0 or 1;

25

or an optical isomer or pharmaceutically acceptable salt thereof.

In a further embodiment of this invention, the inhibitors of farnesyl-protein transferase are illustrated by the formula G:




wherein:

5 R<sup>1a</sup> and R<sup>1c</sup> are independently selected from: hydrogen, R<sup>8</sup>O-,  
-N(R<sup>8</sup>)<sub>2</sub>, F, C<sub>3</sub>-C<sub>10</sub> cycloalkyl or C<sub>1</sub>-C<sub>6</sub> alkyl;

R<sup>1b</sup> is independently selected from:

- 10 a) hydrogen,  
b) aryl, heterocycle or C<sub>3</sub>-C<sub>10</sub> cycloalkyl,  
c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl,  
heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, R<sup>8</sup>O-, or  
-N(R<sup>8</sup>)<sub>2</sub>;

15 R<sup>2a</sup> and R<sup>2b'</sup> are independently selected from selected from: H;  
C<sub>1</sub>-C<sub>6</sub> alkyl,



R<sup>3a</sup> and R<sup>3b</sup> are independently selected from:

- 20 a) hydrogen,  
b) unsubstituted or substituted aryl, unsubstituted or  
substituted heterocycle, unsubstituted or substituted C<sub>3</sub>-C<sub>10</sub>  
cycloalkyl, unsubstituted or substituted C<sub>2</sub>-C<sub>6</sub> alkenyl,  
C<sub>2</sub>-C<sub>6</sub> alkynyl, halogen, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>9</sup>O-,  
R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>9</sup>C(O)O-,

$R^8_2N-C(NR^8)-$ , CN,  $NO_2$ ,  $R^8C(O)-$ ,  $N_3$ ,  $-N(R^8)_2$ , or  $R^9OC(O)NR^8-$ ,

c) unsubstituted  $C_1-C_6$  alkyl,

d) substituted  $C_1-C_6$  alkyl wherein the substituent on the substituted  $C_1-C_6$  alkyl is selected from unsubstituted or substituted aryl, unsubstituted or substituted heterocyclic,  $C_3-C_{10}$  cycloalkyl,  $C_2-C_6$  alkenyl,  $C_2-C_6$  alkynyl,  $R^9O-$ ,  $R^9S(O)_m-$ ,  $R^8C(O)NR^8-$ ,  $(R^8)_2NC(O)-$ ,  $R^8_2N-C(NR^8)-$ , CN,  $R^8C(O)-$ ,  $N_3$ ,  $-N(R^8)_2$ , and  $R^9OC(O)-NR^8-$ ;

$R^{5a}$  and  $R^{5b}$  are independently hydrogen, ethyl, cyclopropyl or methyl;

$R^6$  and  $R^7$  are independently selected from:

H;  $C_1-4$  alkyl,  $C_3-6$  cycloalkyl, aryl, heterocycle,

unsubstituted or substituted with one or two:

a)  $C_1-4$  alkoxy,

b) halogen, or

c) substituted or unsubstituted aryl or substituted or unsubstituted heterocycle;

$R^8$  is independently selected from hydrogen,  $C_1-C_6$  alkyl, 2,2,2-trifluoroethyl, benzyl and aryl;

$R^9$  is independently selected from  $C_1-C_6$  alkyl and aryl;

J, K and L are independently selected from: S, N, O, NH and CH;

$A^1$  is selected from: a bond,  $-C(O)-$ , O,  $-N(R^8)-$ , or  $-S(O)_m$ ;

X is a bond,  $-S(O)_m-$ , O or  $-C(=O)-$ ;

m is 0, 1 or 2;

n is 0, 1 or 2; provided that n is not 0 if  $A^1$  is a bond, O,

-N(R<sup>8</sup>)-, or S(O)<sub>m</sub>;

p is 1, 2 or 3;

q is 0, 1 or 2, provided that q is not 0 or 1 if X is O; and

u is independently 0 or 1;


5

or an optical isomer or pharmaceutically acceptable salt thereof.

The preferred compounds of this invention are as follows:


10

5-(1-(4-Cyanobenzyl)-5-imidazolylmethyl)-4,5,6,7-tetrahydrothieno[3,2-c]pyridine




15

2-Bromo-5-(1-(4-cyanobenzyl)-5-imidazolylmethyl)-4,5,6,7-tetrahydrothieno[3,2-c]pyridine




20

2-Phenyl-5-(1-(4-cyanobenzyl)-5-imidazolylmethyl)-4,5,6,7-tetrahydrothieno[3,2-c]pyridine




25

5-[5-(5-Azabenzimidazolyl)methyl]-1-(4-cyanobenzyl)imidazole



5 5-[4-(4-Azabenzimidazolyl)methyl]-1-(4-cyanobenzyl)imidazole



or a pharmaceutically acceptable salt thereof.

The compounds of the present invention may have  
 10 asymmetric centers and occur as racemates, racemic mixtures, and as individual diastereomers, with all possible isomers, including optical isomers, being included in the present invention. When any variable (e.g. aryl, heterocycle, R<sup>1a</sup>, R<sup>4</sup> etc.) occurs more than one time in any constituent, its definition on each occurrence is independent at every  
 15 other occurrence. Also, combinations of substituents/or variables are permissible only if such combinations result in stable compounds.


As used herein, "alkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms; "alkoxy" represents an alkyl group  
 20 of indicated number of carbon atoms attached through an oxygen bridge. "Halogen" or "halo" as used herein means fluoro, chloro, bromo and iodo.

As used herein, "aryl" is intended to mean any stable monocyclic, bicyclic or tricyclic carbon ring of up to 7 members in each ring, wherein at least one ring is aromatic. Examples of monocyclic and bicyclic aryl elements include phenyl, naphthyl, tetrahydronaphthyl, indanyl, biphenyl, phenanthryl, anthryl or acenaphthyl. Examples of tricyclic aryl elements include 10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-yl (which is also known as dibenzylsuberyl), 9-fluorenyl and 9,10-dihydroanthracen-9-yl. Preferably, "aryl" is a monocyclic or bicyclic carbon ring.

The term heterocycle or heterocyclic, as used herein, represents a stable 5- to 7-membered monocyclic or stable 8- to 11-membered bicyclic heterocyclic ring or stable 13- to 15-membered tricyclic heterocyclic ring, which is either saturated or unsaturated, and which consists of carbon atoms and from one to four heteroatoms selected from the group consisting of N, O, and S, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The heterocyclic ring may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. Examples of monocyclic and bicyclic heterocyclic elements include, but are not limited to, azepinyl, benzimidazolyl, benzisoxazolyl, benzofurazanyl, benzopyranyl, benzothiopyranyl, benzofuryl, benzothiazolyl, benzothienyl, benzoxazolyl, chromanyl, cinnoliny, dihydrobenzofuryl, dihydrobenzothienyl, dihydrobenzothiopyranyl, dihydrobenzothiopyranyl sulfone, furyl, imidazolidinyl, imidazolinyl, imidazolyl, indoliny, indolyl, isochromanyl, isoindoliny, isoquinoliny, isothiazolidinyl, isothiazolyl, isothiazolidinyl, morpholiny, naphthyridinyl, oxadiazolyl, 2-oxoazepinyl, 2-oxopyrrolidinyl, pyridyl, pyrazinyl, pyrazolidinyl, pyrazolyl, pyridazinyl, pyrimidinyl, pyrrolidinyl, pyrrolyl, quinazolinyl, quinoliny, quinoxaliny, tetrahydrofuryl, tetrahydroisoquinoliny, tetrahydroquinoliny, thiamorpholiny, thiamorpholiny sulfoxide, thiazolyl, thiazolinyl, thienofuryl, thienothienyl, and thienyl. Examples of tricyclic heterocyclic elements include, but are not limited to, 6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine, 9,10-dihydro-4H-3-thia-



benzo[f]azulen-4-yl and 9-xanthenyl. The 6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine moiety has the following structure:



- 5 Preferably, "heterocyclic" is a monocyclic or bicyclic moiety.
- As used herein, "heteroaryl" is intended to mean any stable monocyclic, bicyclic or tricyclic carbon ring of up to 7 members in each ring, wherein at least one ring is aromatic and wherein from one to four carbon atoms are replaced by heteroatoms selected from the group
- 10 consisting of N, O, and S. Examples of monocyclic and bicyclic heteroaryl elements include, but are not limited to, benzimidazolyl, benzisoxazolyl, benzofurazanyl, benzopyranyl, benzothiopyranyl, benzofuryl, benzothiazolyl, benzothienyl, benzoxazolyl, chromanyl, cinnolinyl, dihydrobenzofuryl, dihydrobenzothienyl, dihydrobenzo-
- 15 thiopyranyl, dihydrobenzothiopyranyl sulfone, furyl, imidazolyl, indolinyl, indolyl, isochromanyl, isoindolinyl, isoquinolinyl, isothiazolyl, naphthyridinyl, oxadiazolyl, pyridyl, pyrazinyl, pyrazolyl, pyridazinyl, pyrimidinyl, pyrrolyl, quinazolinyl, quinolinyl, quinoxalinyl, tetrahydroisoquinolinyl, tetrahydroquinolinyl, thiazolyl,
- 20 thienofuryl, thienothienyl, and thienyl. Examples of tricyclic heteroaryl elements include, but are not limited to, 6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine. Preferably, "heteroaryl" is a monocyclic or bicyclic moiety.


- As used herein, the terms "substituted aryl", "substituted
- 25 heterocycle" and "substituted cycloalkyl" are intended to include the cyclic group containing from 1 to 3 substituents in addition to the point of attachment to the rest of the compound. Such substituents are preferably selected from the group which includes but is not limited to F, Cl, Br, CF<sub>3</sub>, NH<sub>2</sub>, N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, NO<sub>2</sub>, CN, (C<sub>1</sub>-C<sub>6</sub> alkyl)O-, -
- 30 OH, (C<sub>1</sub>-C<sub>6</sub> alkyl)S(O)<sub>m</sub>-, (C<sub>1</sub>-C<sub>6</sub> alkyl)C(O)NH-, H<sub>2</sub>N-C(NH)-, (C<sub>1</sub>-

C<sub>6</sub> alkyl)C(O)-, (C<sub>1</sub>-C<sub>6</sub> alkyl)OC(O)-, N<sub>3</sub>-(C<sub>1</sub>-C<sub>6</sub> alkyl)OC(O)NH- and C<sub>1</sub>-C<sub>20</sub> alkyl.


When R<sup>6</sup> and R<sup>7</sup> are combined to form a ring, cyclic amine moieties are formed. Examples of such cyclic moieties include,  
 5 but are not limited to:


















In addition, such cyclic moieties may optionally include another heteroatom(s). Examples of such heteroatom-containing cyclic amine moieties include, but are not limited to:






















The variable J, K and L are selected such that the 5-member ring comprising them is aromatic in nature. Examples of the  
 15 following ring system




include, but are not limited to:









Preferably the ring system is selected from:



## 5 Examples of the following ring system




include, but are not limited to:



Lines drawn into the ring systems from substituents (such as from R<sup>3a</sup>, R<sup>3b</sup>, R<sup>4</sup> etc.) indicate that the indicated bond may be attached to any of the substitutable ring carbon or nitrogen atoms.

Preferably, R<sup>1a</sup> and R<sup>1b</sup> are independently selected from: hydrogen, -N(R<sup>8</sup>)<sub>2</sub>, R<sup>8</sup>C(O)NR<sup>8</sup>- or C<sub>1</sub>-C<sub>6</sub> alkyl which is unsubstituted or substituted by -N(R<sup>8</sup>)<sub>2</sub>, R<sup>8</sup>O- or R<sup>8</sup>C(O)NR<sup>8</sup>-.

Preferably, R<sup>2a</sup> is selected from: H;



Preferably, R<sup>2b'</sup> and R<sup>2b''</sup> are independently selected from selected from: H or C<sub>1</sub>-C<sub>6</sub> alkyl.

Preferably, R<sup>3a</sup> and R<sup>3b</sup> are independently selected from: hydrogen, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, CN, R<sup>8</sup>C(O)-, -N(R<sup>8</sup>)<sub>2</sub> and C<sub>1</sub>-C<sub>6</sub> alkyl.

Preferably, R<sup>4</sup> is selected from: hydrogen, perfluoroalkyl, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, CN, NO<sub>2</sub>, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, R<sup>9</sup>OC(O)NR<sup>8</sup>- and C<sub>1</sub>-C<sub>6</sub> alkyl.

Preferably, R<sup>5</sup> is hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl.

Preferably, R<sup>8</sup> is selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl and benzyl.

Preferably, A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond, -C(O)NR<sup>8</sup>-, -NR<sup>8</sup>C(O)-, O, -N(R<sup>8</sup>)-, -S(O)<sub>2</sub>N(R<sup>8</sup>)- and -N(R<sup>8</sup>)S(O)<sub>2</sub>-.

Preferably, V is selected from hydrogen, heterocycle and aryl.

Preferably, W is imidazolyl.

Preferably, X is a bond or (C=O)-.

Preferably, n, p and r are independently 0, 1, or 2.


More preferably, r is 1.

Preferably t is 1.


Preferably u is independently 0 or 1. Most preferably,

5 u is 1.

Preferably, the moiety



is selected from:



10

The pharmaceutically acceptable salts of the compounds of this invention include the conventional non-toxic salts of the compounds of this invention as formed, e.g., from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those  
 15 derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxy-benzoic,  
 20 fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, trifluoroacetic and the like.

It is intended that the definition of any substituent or variable (e.g., R<sup>1a</sup>, Z, n, etc.) at a particular location in a molecule be independent of its definitions elsewhere in that molecule. Thus,  
 25 -N(R<sup>8</sup>)<sub>2</sub> represents -NH<sub>2</sub>, -NHCH<sub>3</sub>, -NHC<sub>2</sub>H<sub>5</sub>, etc. It is understood that substituents and substitution patterns on the compounds of the

instant invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art, as well as those methods set forth below, from readily available starting materials.

- 5           The pharmaceutically acceptable salts of the compounds of this invention can be synthesized from the compounds of this invention which contain a basic moiety by conventional chemical methods. Generally, the salts are prepared either by ion exchange chromatography or by reacting the free base with stoichiometric  
10 amounts or with an excess of the desired salt-forming inorganic or organic acid in a suitable solvent or various combinations of solvents.

Abbreviations used in the description of the chemistry and in the Examples that follow are:

15	Ac <sub>2</sub> O	Acetic anhydride;
	Boc	t-Butoxycarbonyl;
	CBz	Carbobenzyloxy;
	DBU	1,8-diazabicyclo[5.4.0]undec-7-ene;
	DMAP	4-Dimethylaminopyridine;
20	DME	1,2-Dimethoxyethane;
	DMF	Dimethylformamide;
	EDC	1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide- hydrochloride;
	Et <sub>3</sub> N	Triethylamine;
25	EtOAc	Ethyl acetate;
	FAB	Fast atom bombardment;
	HOBT	1-Hydroxybenzotriazole hydrate;
	HOBT	3-Hydroxy-1,2,2-benzotriazin-4(3 <i>H</i> )-one;
	HPLC	High-performance liquid chromatography;
30	MCPBA	m-Chloroperoxybenzoic acid;
	MsCl	Methanesulfonyl chloride;
	NaHMDS	Sodium bis(trimethylsilyl)amide;
	Py	Pyridine;
	TFA	Trifluoroacetic acid;



THF            Tetrahydrofuran.

5            The compounds of this invention are prepared by  
employing reactions as shown in the Schemes 1-16, in addition to other  
standard manipulations such as ester hydrolysis, cleavage of protecting  
groups, etc., as may be known in the literature or exemplified in the  
experimental procedures. While stereochemistry is shown in the  
Schemes, a person of ordinary skill in the art would understand that the  
10            illustrated compounds represent racemic mixtures which may be  
separated at a subsequent purification step or may be utilized as the  
racemic mixture.

15            These reactions may be employed in a linear sequence to  
provide the compounds of the invention or they may be used to  
synthesize fragments which are subsequently joined by the reductive  
alkylation or acylation reactions described in the Schemes.

Synopsis of Schemes 1-7:

20            The requisite intermediates are in some cases commercially  
available, or can be prepared according to literature procedures, for the  
most part. For example, see M. Cain *et al.*, *Heterocycles*, 19:1003  
(1982), J.W. Skiles *et al.*, *J. Med. Chem.*, 29:784 (1986) and J.L.  
Stanton *et al.*, *J. Med. Chem.*, 26:1267 (1983).

25            In Schemes 1-2, for example, the syntheses of several fused  
ring intermediates are outlined. The subsequent reactions described in  
the remaining schemes may be similarly applied to suitably protected  
commercially available tetrahydroisoquinolines, as well as commercially  
or synthetically obtained homologs, to provide compounds of the instant  
invention.

30            Scheme 1 describes the synthesis of 4,5,6,7-  
tetrahydrothieno[3,2-c]pyridines essentially according to the method of  
Halczenko and Hartman in *Synthetic Commun.*, **1996**, 26, 1363. The  
nitrile **1** may be reduced to the amine **2** using, for example, Raney  
Nickel and this compound undergoes cyclization to **3** when heated with  
formaldehyde followed by hydrochloric acid treatment. Intermediate **3**

may then be functionalized as described in subsequent schemes. The regioisomeric thienopyridine **4** is known (Gronowitz, *Ark. Kemi.*; **1970**, 32, 217) and may undergo the same reactions as are described for **3**.

5                    Synthesis of the lower homolog of **3** is shown in Scheme 2. Using a modification of the chemistry described in US patent 5,334,596, the commercially available 2,3-thiophenedicarboxylic acid anhydride may be heated with an amine to yield **5**. Reduction of the carbonyls using, for example, diborane in a solvent such as THF would yield  
10                    analogs such as **6**. Further derivatives could be prepared by first deprotecting a suitably functionalized **6**.

                    The furano analogs **7** (Scheme 2a) are described in the literature (Arnoldi et al, *J. Heterocyclic Chem.*, **1990**, 27, 1169). Deprotection using, for example, aqueous HCl will produce **8** which  
15                    may undergo reactions as described hereinbelow.

                    Scheme 3 illustrates reactions wherein the preferred 4-cyanobenzylimidazolyl moiety is incorporated into the instant compounds, either by reductive alkylation with the aldehyde **9** or an amide forming reaction with the acid **10**.

20                    Schemes 4-5 illustrate the syntheses of 4,5,6,7-tetrahydrothieno[3,2-c]pyridine of the instant invention wherein the "X" moiety is other than an alkyl bridge. The reactions illustrated therein show the incorporation of sidechains which comprise the preferred 4-cyanobenzylimidazolyl moiety. It is understood that a person of  
25                    ordinary skill in the art could readily modify such reaction sequences by using appropriate protecting groups and reagents well known to one skilled in the art to provide other compounds of the instant invention.


                    Scheme 4 illustrates the syntheses of compounds of the instant invention wherein "X" is -S-. Thus the intermediate aldehyde **11**  
30                    is reduced to the alcohol **12**, activated and treated with a suitable thioacetate to provide the thioester **13**. The thiol is then generated and alkylated with a suitable ester containing reagent, such as bromoacetic acid to provide intermediate **14**. Reduction of the ester moiety, followed by oxidation provides the corresponding aldehyde, which can


be utilized to reductively alkylate the suitably substituted 4,5,6,7-tetrahydrothieno[3,2-c]pyridine, for example, to provide the instant compound **15**.

Scheme 5 illustrates the syntheses of compounds of the instant invention wherein "X" is -O-. Thus, a dihydroxyalkane, such as ethylene glycol, can be selectively protected and oxidized to provide the aldehyde **16**. Intermediate **16** can be utilized to reductively alkylate the suitably substituted of 4,5,6,7-tetrahydro-thieno[3,2-c]pyridine, the sidechain deprotected and converted to the chloride **17**. Intermediate **17** can then be alkylated with a suitable reagent to provide the instant compound **18** which incorporates the ether moiety.


The reagent utilized in the reductive alkylation of the of 4,5,6,7-tetrahydrothieno[3,2-c]pyridine may alternatively incorporate a leaving group which may subsequently react with a blocked imidazolyl reagent, such as **19** to provide compounds of the instant invention wherein "X" is a bond and the preferred imidazolyl is attached to the alkyl bridge via one of the ring nitrogens, as shown in Scheme 6.


Scheme 7 illustrates the syntheses of compounds of the instant invention comprising 3,4-dihydro-1(1H)-thieno[3,2-c]pyridones and a homologous [3.3.0] ring system.

SCHEME 1


SCHEME 2


10

SCHEME 2a


SCHEME 3


## SCHEME 4




SCHEME 4 (continued)



SCHEME 5

SCHEME 6

SCHEME 7

It is understood that while Schemes 8-16 illustrate preparation of both protected and unprotected intermediates, a person of ordinary skill would appreciate that subsequent reactions which utilize those intermediates, such as those described in Schemes 1-7, may require protection and eventual deprotection of certain intermediate moieties.

The selectively protected intermediate **20** utilized in the synthesis illustrated in Scheme 8 can be reductively alkylated with a variety of aldehydes, such as **21**. The aldehydes can be prepared by standard procedures, such as that described by O. P. Goel, U. Krolls, M. Stier and S. Kesten in Organic Syntheses, **1988**, 67, 69-75. The reductive alkylation can be accomplished at pH 5-7 with a variety of reducing agents, such as sodium triacetoxyborohydride or sodium cyanoborohydride in a solvent such as dichloroethane, methanol or dimethylformamide. The ester product **22** can be deprotected with trifluoroacetic acid in methylene chloride to give the substituted diamine **23**. That diamine may be isolated in the salt form, for example, as a trifluoroacetate, hydrochloride or acetate salt, among others. The product diamine **23** can be further selectively protected and reductively alkylated with a second aldehyde to obtain an analogous tertiary amine. Alternatively, the diamine **23** can be cyclized to obtain intermediates such as the dihydroimidazole **24** by procedures known in the literature. The ester **24** can then be utilized in a reaction such as illustrated in Scheme 3 hereinabove.

Scheme 9 illustrates a general preparation of aralkyl imidazolyl intermediates **31** that can be utilized in reactions such as illustrated in Scheme 3. Thus imidazole acetic acid **27** can be converted to the protected acetate **29** by standard procedures, and **29** can be first reacted with an alkyl halide, then treated with refluxing methanol to provide the regiospecifically alkylated imidazole acetic acid ester **30**. Hydrolysis provides the acetic acid **31**.


Schemes 10-13 illustrate syntheses of suitably substituted alkanols useful in the syntheses of the instant compounds wherein

the variable W is present as a pyridyl moiety. The hydroxyl moiety of such intermediates may be converted into the corresponding aldehyde, as illustrated in Scheme 10 or may be converted to a suitable leaving group, as illustrated in Scheme 12. Similar synthetic strategies for preparing alkanols that incorporate other heterocyclic moieties for variable W are also well known in the art.


Compounds of the instant invention wherein the  $A^1(CR^{1a}_2)_nA^2(CR^{1a}_2)_n$  linker is a substituted methylene may be synthesized by the methods shown in Scheme 14. Thus, the N-protected imidazolyl iodide **32** is reacted, under Grignard conditions with a suitably protected benzaldehyde to provide the alcohol **33**. Acylation, followed by the alkylation procedure illustrated in the Schemes above (in particular, Scheme 6) provides the instant compound **34**. If other  $R^{1a}$  substituents are desired, the acetyl moiety can be manipulated as illustrated in the Scheme.

Scheme 15 illustrates synthesis of an instant compound wherein a non-hydrogen  $R^{5b}$  is incorporated in the instant compound. Thus, a readily available 4-substituted imidazole **37** may be selectively iodinated to provide the 5-iodoimidazole **38**. That imidazole may then be protected and coupled to a suitably substituted benzyl moiety to provide intermediate **39**. Intermediate **39** can then undergo the alkylation reactions that were described hereinabove.

Compounds of the instant invention wherein the  $A^1(CR^{1a}_2)_nA^2(CR^{1a}_2)_n$  linker is oxygen may be synthesized by methods known in the art, for example as shown in Scheme 16. The suitably substituted phenol **41** may be reacted with methyl N-(cyano)methanimidate to provide the 4-phenoxyimidazole **42**. After selective protection of one of the imidazolyl nitrogens, the intermediate **43** can undergo alkylation reactions as described for the benzylimidazoles hereinabove.


SCHEME 8


## SCHEME 8 (continued)



5


## SCHEME 9





SCHEME 10




## SCHEME 11





SCHEME 12


SCHEME 13

## SCHEME 14




SCHEME 14 (continued)

## SCHEME 15



## SCHEME 16



The instant compounds are useful as pharmaceutical agents for mammals, especially for humans. These compounds may be administered to patients for use in the treatment of cancer. Examples of the type of cancer which may be treated with the compounds of this invention include, but are not limited to, colorectal carcinoma, exocrine pancreatic carcinoma, myeloid leukemias and neurological tumors. Such tumors may arise by mutations in the *ras* genes themselves, mutations in the proteins that can regulate Ras activity (i.e., neurofibromin (NF-1), neu, scr, abl, lck, fyn) or by other mechanisms.

The compounds of the instant invention inhibit farnesyl-protein transferase and the farnesylation of the oncogene protein Ras. The instant compounds may also inhibit tumor angiogenesis, thereby affecting the growth of tumors (J. Rak et al. *Cancer Research*, 55:4575-4580 (1995)). Such anti-angiogenesis properties of the instant compounds may also be useful in the treatment of certain forms of blindness related to retinal vascularization.

The compounds of this invention are also useful for inhibiting other proliferative diseases, both benign and malignant, wherein Ras proteins are aberrantly activated as a result of oncogenic mutation in other genes (i.e., the Ras gene itself is not activated by mutation to an oncogenic form) with said inhibition being accomplished by the administration of an effective amount of the compounds of the invention to a mammal in need of such treatment. For example, the compounds are useful in the treatment of neurofibromatosis, which is a benign proliferative disorder.

The instant compounds may also be useful in the treatment of certain viral infections, in particular in the treatment of hepatitis delta and related viruses (J.S. Glenn et al. *Science*, 256:1331-1333 (1992)).

The compounds of the instant invention are also useful in the prevention of restenosis after percutaneous transluminal coronary angioplasty by inhibiting neointimal formation (C. Indolfi et al. *Nature medicine*, 1:541-545(1995)).



The instant compounds may also be useful in the treatment and prevention of polycystic kidney disease (D.L. Schaffner et al. *American Journal of Pathology*, 142:1051-1060 (1993) and B. Cowley, Jr. et al. *FASEB Journal*, 2:A3160 (1988)).

5           The instant compounds may also be useful for the treatment of fungal infections.

          In a preferred embodiment of the instant invention the compounds of this instant invention are selective inhibitors of farnesyl-protein transferase. A compound is considered a selective inhibitor of  
10   farnesyl-protein transferase, for example, when its *in vitro* farnesyl-protein transferase inhibitory activity, as assessed by the assay described in Example 6, is at least 100 times greater than the *in vitro* activity of the same compound against geranylgeranyl-protein transferase-type I in the assay described in Example 7. Preferably, a selective compound  
15   exhibits at least 1000 times greater activity against one of the enzymatic activities when comparing geranylgeranyl-protein transferase-type I inhibition and farnesyl-protein transferase inhibition.

          In another preferred embodiment of the instant invention the compounds of this instant invention are dual inhibitors of farnesyl-protein transferase and geranylgeranyl-protein transferase type I. Such a  
20   dual inhibitor will exhibit certain characteristics when assessed in *in vitro* assays, which are dependent on the type of assay employed.

          In a SEAP assay, such as described in Example 10, it is preferred that the dual inhibitor compound has an *in vitro* inhibitory  
25   activity (IC<sub>50</sub>) that is less than about 12μM against K4B-Ras dependent activation of MAP kinases in cells. More preferably, the dual inhibitor compound has an *in vitro* inhibitory activity (IC<sub>50</sub>) against K4B-Ras dependent activation of MAP kinases in cells which is more than about 5 times lower than the inhibitory activity (IC<sub>50</sub>) against Myr-Ras  
30   dependent activation of MAP kinases in cells. Also more preferably, in a SEAP assay, the dual inhibitor compound has an inhibitory activity (IC<sub>50</sub>) that is less than about 10 nM against H-Ras dependent activation of MAP kinases in cells.

In a GGTase plus anion assay, such as described in Example 7, it is preferred that the dual inhibitor compound has an *in vitro* inhibitory activity (IC<sub>50</sub>) that is less than about 5  $\mu$ M against transfer of a geranylgeranyl residue to a protein or peptide substrate comprising a CAAX<sup>G</sup> motif by geranylgeranyl-protein transferase type I in the presence of a modulating anion. More preferably, the dual inhibitor compound has an *in vitro* inhibitory activity (IC<sub>50</sub>) that is less than about 1  $\mu$ M against transfer of a geranylgeranyl residue to a protein or peptide substrate comprising a CAAX<sup>G</sup> motif by geranylgeranyl-protein transferase type I in the presence of a modulating anion. Preferably, the dual inhibitor compound has an *in vitro* inhibitory activity (IC<sub>50</sub>) in the *in vitro* assay as described in Example 6 that is less than about 1  $\mu$ M against transfer of a farnesyl residue to a protein or peptide substrate, comprising a CAAX<sup>F</sup> motif, by farnesyl-protein transferase. more preferably, the dual inhibitor compound has an *in vitro* inhibitory activity (IC<sub>50</sub>) that is less than about 100nM against transfer of a farnesyl residue to a protein or peptide substrate, comprising a CAAX<sup>F</sup> motif, by farnesyl-protein transferase. Also preferably, the dual inhibitor compound has an *in vitro* inhibitory activity (IC<sub>50</sub>) in the *in vitro* assay as described in Example 9, that is less than about 100 nM against the anchorage independent growth of H-*ras*-transformed mammalian fibroblasts.

The protein or peptide substrate utilized in the instant assay may incorporate any CAAX motif that is geranylgeranylated by GGTase-I. The term "CAAX<sup>G</sup>" will refer to such motifs that may be geranylgeranylated by GGTase-I. It is understood that some of the "CAAX<sup>G</sup>" containing protein or peptide substrates may also be farnesylated by farnesyl-protein transferase. In particular such "CAAX<sup>G</sup>" motifs include (the corresponding human protein is in parentheses): CVIM (K4B-Ras) (SEQ.ID.NO.: 1), CVLL (mutated H-Ras) (SEQ.ID.NO.: 2), CVVM (N-Ras) (SEQ.ID.NO.: 3), CIIM (K4A-Ras) (SEQ.ID.NO.: 4), CLLL (Rap-1A) (SEQ.ID.NO.: 5), CQLL (Rap-1B) (SEQ.ID.NO.: 6), CSIM (SEQ.ID.NO.: 7), CAIM (SEQ.ID.NO.: 8),

CKVL (SEQ.ID.NO.: 9) and CLIM (PFX) (SEQ.ID.NO.: 10).

Preferably, the CAAX motif is CVIM (SEQ.ID.NO.: 1).

As used herein, the term "CAAX<sup>F</sup>" is used to designate a protein or peptide substrate that incorporates four amino acid C-terminus motif that is farnesylated by farnesyl-protein transferase. It is understood that certain of the "CAAX<sup>F</sup>" containing protein or peptide substrates may also be geranylgeranylated by GGTase-I. In particular such "CAAX<sup>F</sup>" motifs include (the corresponding human protein is in parentheses): CVLS (H-ras) (SEQ.ID.NO.: 11), CVIM (K4B-Ras) (SEQ.ID.NO.: 1) and CVVM (N-Ras) (SEQ.ID.NO.: 3).

The compounds of this invention may be administered to mammals, preferably humans, either alone or, preferably, in combination with pharmaceutically acceptable carriers or diluents, optionally with known adjuvants, such as alum, in a pharmaceutical composition, according to standard pharmaceutical practice. The compounds can be administered orally or parenterally, including the intravenous, intramuscular, intraperitoneal, subcutaneous, rectal and topical routes of administration.

For oral use of a chemotherapeutic compound according to this invention, the selected compound may be administered, for example, in the form of tablets or capsules, or as an aqueous solution or suspension. In the case of tablets for oral use, carriers which are commonly used include lactose and corn starch, and lubricating agents, such as magnesium stearate, are commonly added. For oral administration in capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening and/or flavoring agents may be added. For intramuscular, intraperitoneal, subcutaneous and intravenous use, sterile solutions of the active ingredient are usually prepared, and the pH of the solutions should be suitably adjusted and buffered. For intravenous use, the total concentration of solutes should be controlled in order to render the preparation isotonic.

The present invention also encompasses a pharmaceutical composition useful in the treatment of cancer, comprising the administration of a therapeutically effective amount of the compounds of this invention, with or without pharmaceutically acceptable carriers or diluents. Suitable compositions of this invention include aqueous solutions comprising compounds of this invention and pharmacologically acceptable carriers, e.g., saline, at a pH level, e.g., 7.4. The solutions may be introduced into a patient's intramuscular blood-stream by local bolus injection.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specific amounts, as well as any product which results, directly or indirectly, from combination of the specific ingredients in the specified amounts.

When a compound according to this invention is administered into a human subject, the daily dosage will normally be determined by the prescribing physician with the dosage generally varying according to the age, weight, and response of the individual patient, as well as the severity of the patient's symptoms.

In one exemplary application, a suitable amount of compound is administered to a mammal undergoing treatment for cancer. Administration occurs in an amount between about 0.1 mg/kg of body weight to about 60 mg/kg of body weight per day, preferably of between 0.5 mg/kg of body weight to about 40 mg/kg of body weight per day.


The compounds of the instant invention are also useful as a component in an assay to rapidly determine the presence and quantity of farnesyl-protein transferase (FPTase) in a composition. Thus the composition to be tested may be divided and the two portions contacted with mixtures which comprise a known substrate of FPTase (for example a tetrapeptide having a cysteine at the amine terminus) and farnesyl pyrophosphate and, in one of the mixtures, a compound of the instant invention. After the assay mixtures are incubated for an sufficient period of time, well known in the art, to

- allow the FPTase to farnesylate the substrate, the chemical content of the assay mixtures may be determined by well known immunological, radiochemical or chromatographic techniques. Because the compounds of the instant invention are selective inhibitors of
- 5 FPTase, absence or quantitative reduction of the amount of substrate in the assay mixture without the compound of the instant invention relative to the presence of the unchanged substrate in the assay containing the instant compound is indicative of the presence of FPTase in the composition to be tested.
- 10 It would be readily apparent to one of ordinary skill in the art that such an assay as described above would be useful in identifying tissue samples which contain farnesyl-protein transferase and quantitating the enzyme. Thus, potent inhibitor compounds of the instant invention may be used in an active site titration assay to determine
- 15 the quantity of enzyme in the sample. A series of samples composed of aliquots of a tissue extract containing an unknown amount of farnesyl-protein transferase, an excess amount of a known substrate of FPTase (for example a tetrapeptide having a cysteine at the amine terminus) and farnesyl pyrophosphate are incubated for an
- 20 appropriate period of time in the presence of varying concentrations of a compound of the instant invention. The concentration of a sufficiently potent inhibitor (i.e., one that has a  $K_i$  substantially smaller than the concentration of enzyme in the assay vessel) required to inhibit the enzymatic activity of the sample by 50% is
- 25 approximately equal to half of the concentration of the enzyme in that particular sample.

### EXAMPLES

- 30 Examples provided are intended to assist in a further understanding of the invention. Particular materials employed, species and conditions are intended to be further illustrative of

the invention and not limitative of the reasonable scope thereof. Purification by HPLC was utilized for the Examples as set forth below.

EXAMPLE 1

5    5-(1-(4-Cyanobenzyl)-5-imidazolylmethyl)-4,5,6,7-  
tetrahydrothieno[3,2-c]pyridine

---

10    Step 1:    Preparation of 1-triphenylmethyl-4-(hydroxymethyl)-  
imidazole

---

To a solution of 4-(hydroxymethyl)imidazole hydrochloride (35 g) in 250 mL of dry DMF at room temperature was added triethylamine (90.6 mL). A white solid precipitated from the solution. Chlorotriphenylmethane (76.1 g) in 500 mL of DMF was added dropwise. The reaction mixture was stirred for 20 hours, poured over ice, filtered, and washed with ice water. The resulting product was slurried with cold dioxane, filtered, and dried *in vacuo* to provide the title compound as a white solid which was sufficiently pure for use in the next step.

20    Step 2:    Preparation of 1-triphenylmethyl-4-(acetoxymethyl)-  
imidazole

---

The alcohol prepared above was suspended in 500 mL of pyridine. Acetic anhydride (74 mL) was added dropwise, and the reaction was stirred for 48 hours during which it became homogeneous. The solution was poured into 2 L of EtOAc, washed with water (3 x 1 L), 5% aq. HCl soln. (2 x 1 L), sat. aq. NaHCO<sub>3</sub>, and brine, then dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated *in vacuo* to provide the crude product. The title compound was isolated as a white powder which was sufficiently pure for use in the next step.

30

Step 3: Preparation of 1-(4-cyanobenzyl)-5-(acetoxymethyl)-  
imidazole hydrobromide

---

A solution of the acetate from Step 2 (85.8 g) and  $\alpha$ -  
bromo-*p*-tolunitrile (50.1 g) in 500 mL of EtOAc was stirred at 60  
5 °C for 20 hours, during which a pale yellow precipitate formed. The  
reaction was cooled to room temperature and filtered to provide the  
solid imidazolium bromide salt. The filtrate was concentrated *in*  
*vacuo* to a volume 200 mL, reheated at 60 °C for two hours, cooled  
to room temperature, and filtered again. The filtrate was concen-  
10 trated *in vacuo* to a volume 100 mL, reheated at 60 °C for another  
two hours, cooled to room temperature, and concentrated *in vacuo* to  
provide a pale yellow solid. All of the solid material was combined,  
dissolved in 500 mL of methanol, and warmed to 60 °C. After two  
hours, the solution was reconcentrated *in vacuo* to provide a white  
15 solid which was triturated with hexane to remove soluble materials.  
Removal of residual solvents *in vacuo* provided the title  
hydrobromide as a white solid which was used in the next step  
without further purification.

20 Step 4: Preparation of 1-(4-cyanobenzyl)-5-(hydroxymethyl)-  
imidazole

---

To a solution of the product from Step 3 (50.4 g) in  
1.5 L of 3:1 THF/water at 0 °C was added lithium hydroxide mono-  
hydrate (18.9 g). After one hour, the reaction was concentrated  
25 *in vacuo*, diluted with EtOAc (3 L), and washed with water, sat. aq.  
NaHCO<sub>3</sub> and brine. The solution was then dried (Na<sub>2</sub>SO<sub>4</sub>), filtered,  
and concentrated *in vacuo* to provide the crude product as a pale  
yellow fluffy solid which was sufficiently pure for use in the next  
step without further purification.

30



Step 5: Preparation of 1-(4-cyanobenzyl)-5-imidazole-carboxaldehyde


To a solution of the alcohol from Step 4 (21.5 g) in 500 mL of DMSO at room temperature was added triethylamine (56 mL), then SO<sub>3</sub>-pyridine complex (40.5 g). After 45 minutes, the reaction was poured into 2.5 L of EtOAc, washed with water (4 x 1 L) and brine, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated *in vacuo* to provide the title aldehyde as a white powder which was sufficiently pure for use in the next step without further purification.

Step 6: Preparation of 5-(1-(4-cyanobenzyl)-5-imidazolylmethyl)-4,5,6,7-tetrahydrothieno[3,2-c]pyridine

To a solution of 4,5,6,7-tetrahydrothieno[3,2-c]pyridine hydrochloride (Halczenko and Hartman; *Synthetic Commun.*, 1996, 26, 1363; 0.439 g, 2.5 mmol) in CHCl<sub>2</sub>CHCl<sub>2</sub> (15 mL) was added 1-(4-cyanobenzyl)-5-imidazole carboxaldehyde (0.58 g, 2.75 mmol), 4Å sieves and triethylamine (0.35 mL, 2.5 mmol) followed by NaBH(OAc)<sub>3</sub> (1.59 g, 7.5 mmol). The mixture was stirred at room temperature for 16 h., the mixture was diluted with EtOAc, filtered through celite, washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated. The residue was stirred in 2N HCl (50 mL) for 2 h. then extracted with ether. The aqueous layer was basified with 40% NaOH solution and extracted 3X CH<sub>2</sub>Cl<sub>2</sub> which was then washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvent evaporated in vacuo. Chromatography (silica gel; 2.5% MeOH in CHCl<sub>3</sub>) followed by crystallization from ether afforded the title compound as a white solid.

Analysis for C<sub>19</sub>H<sub>18</sub>N<sub>4</sub>S•0.05 CHCl<sub>3</sub>

	Calcd.	C, 67.21; H, 5.34; N, 16.46
30	found	C, 67.27; H, 5.26; N, 16.61

EXAMPLE 2

5    2-Bromo-5-(1-(4-cyanobenzyl)-5-imidazolylmethyl)-4,5,6,7-tetrahydrothieno[3,2-c]pyridine

Step 1:    Preparation of N-Boc-4,5,6,7-tetrahydrothieno[3,2-c]pyridine

10            4,5,6,7-Tetrahydrothieno[3,2-c]pyridine.HCl (7.38 g, 42 mmol) in CHCl<sub>3</sub> (150 mL) was treated with triethylamine (11.7 mL, 84 mmol) and (Boc)<sub>2</sub>O (10.1 g, 46.2 mmol) for 5 h. The CHCl<sub>3</sub> was removed in vacuo and the residue partitioned between water and EtOAc. Extracted with EtOAc (3X), washed successively with 10%  
15    KHSO<sub>4</sub>, brine, saturated NaHCO<sub>3</sub> then brine, dried and evaporated to give the title compound as a solid.

Step 2:    Preparation of 2-bromo-4,5,6,7-tetrahydrothieno[3,2-c]pyridine hydrobromide


20            The product from Step 1 (9.57 g, 40 mmol) was dissolved in CHCl<sub>3</sub> (150 mL) at 0°C and treated with bromine (2.1 mL, 40 mmol) dropwise over 10 minutes. The mixture was stirred for 16 h at room temperature then filtered to afford the title compound as a solid.

25            Step 3:    Preparation of 2-bromo-5-(1-(4-cyanobenzyl)-5-imidazolylmethyl)-4,5,6,7-tetrahydrothieno[3,2-c]pyridine

30            Following the procedure described for Example 1, Step 6 but using 2-bromo-4,5,6,7-tetrahydrothieno[3,2-c]pyridine the title compound was obtained as a white solid.  
Analysis for C<sub>19</sub>H<sub>17</sub>N<sub>4</sub>BrS

Calcd. C, 55.21; H, 4.15; N, 13.56  
 found C, 55.67; H, 4.26; N, 13.59

### EXAMPLE 3



2-Phenyl-5-(1-(4-cyanobenzyl)-5-imidazolylmethyl)-4,5,6,7-tetrahydrothieno[3,2-c]pyridine

10

Step 1: Preparation of N-Boc-2-bromo-4,5,6,7-tetrahydrothieno[3,2-c]pyridine

2-bromo-4,5,6,7-tetrahydrothieno[3,2-c]pyridine hydrobromide (Example 2, Step 2; 6.67 g, 25 mmol) was converted to the Boc-protected derivative using (Boc)<sub>2</sub>O (5.46 g, 25 mmol) in 150 mL dioxane and 75 mL 1N NaOH for 2 h. Work-up as described for Example 2, Step 1 afforded the title compound as a solid.

20 Step 2: Preparation of N-Boc-2-phenyl-4,5,6,7-tetrahydrothieno[3,2-c]pyridine

A solution of N-Boc-2-bromo-4,5,6,7-tetrahydrothieno[3,2-c]pyridine (Step 1; 637 mg, 0.2 mmol), phenyl boronic acid (244 mg, 0.2 mmol), NaHCO<sub>3</sub> (504 mg, 6 mmol) and tetrakis(triphenylphosphine)palladium(0) (116 mg, 0.1 mmol) in DME (10 mL) and water (10 mL) was heated to 80°C for 20 h. The mixture was partitioned between water and EtOAc, washed with aqueous NaHCO<sub>3</sub>, water (2X) then brine, dried and evaporated. Column chromatography of the residue (silica gel; hexane/EtOAc 95:5) afforded the title compound as a viscous oil.

Step 3: Preparation of 2-phenyl-4,5,6,7-tetrahydrothieno[3,2-c]pyridine hydrochloride

The product from Step 2 was dissolved in EtOAc at -25°C and treated with HCl gas until saturated. the reaction was stoppered and stirred at 0°C for 1h. Removal of the solvent in vacuo afforded the title compound as a solid.


Step 4: Preparation of 2-phenyl-5-(1-(4-cyanobenzyl)-5-imidazolylmethyl)-4,5,6,7-tetrahydrothieno[3,2-c]pyridine

Following the procedure described for Example 1, Step 6 but using 2-phenyl-4,5,6,7-tetrahydrothieno[3,2-c]pyridine.HCl the title compound was obtained as a white solid.

Analysis for  $C_{25}H_{22}N_4S \cdot 0.25H_2O \cdot 0.1Et_2O$

Calcd.	C, 72.21; H, 5.61; N, 13.26
found	C, 72.39; H, 5.31; N, 13.00

EXAMPLE 4



5-[5-(5-Azabenzimidazolyl)methyl]-1-(4-cyanobenzyl)imidazole hydrochloride

Step 1: Preparation of 1-triphenylmethyl-4-(hydroxymethyl)-imidazole

To a solution of 4-(hydroxymethyl)imidazole hydrochloride (35.0 g, 260 mmol) in 250 mL of dry DMF at room temperature was added triethylamine (90.6 mL, 650 mmol). A white solid precipitated from the solution. Chlorotriphenylmethane (76.1 g, 273 mmol) in 500 mL of DMF was added dropwise. The reaction

mixture was stirred for 20 hours, poured over ice, filtered, and washed with ice water. The resulting product was slurried with cold dioxane, filtered, and dried *in vacuo* to provide the titled product as a white solid which was sufficiently pure for use in the next step.

5

Step 2: Preparation of 1-triphenylmethyl-4-(acetoxymethyl)imidazole

The alcohol described in Step 1 (260 mmol, prepared above) was suspended in 500 mL of pyridine. Acetic anhydride (74 mL, 780 mmol) was added dropwise, and the reaction was stirred for 48 hours during which it became homogeneous. The solution was poured into 2 L of EtOAc, washed with water (3 x 1 L), 5% aq. HCl soln. (2 x 1 L), sat. aq. NaHCO<sub>3</sub>, and brine, then dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated *in vacuo* to provide the crude product. The acetate was isolated as a white powder, which was sufficiently pure for use in the next reaction.

Step 3: Preparation of 1-(4-cyanobenzyl)-5-(acetoxymethyl)imidazole hydrobromide

A solution of the product described in Step 2 (85.8 g, 225 mmol) and  $\alpha$ -bromo-*p*-tolunitrile (50.1 g, 232 mmol) in 500 mL of EtOAc was stirred at 60 °C for 20 hours, during which a pale yellow precipitate formed. The reaction was cooled to room temperature and filtered to provide the solid imidazolium bromide salt. The filtrate was concentrated *in vacuo* to a volume 200 mL, reheated at 60 °C for two hours, cooled to room temperature, and filtered again. The filtrate was concentrated *in vacuo* to a volume 100 mL, reheated at 60 °C for another two hours, cooled to room temperature, and concentrated *in vacuo* to provide a pale yellow solid. All of the solid material was combined, dissolved in 500 mL of methanol, and warmed to 60 °C. After two hours, the solution was reconcentrated *in vacuo* to provide a white solid which was triturated with hexane to remove soluble materials. Removal of residual solvents *in vacuo* provided the titled

product hydrobromide as a white solid, which was used in the next step without further purification.

**Step 4:** Preparation of 1-(4-cyanobenzyl)-5-(hydroxymethyl)-imidazole

To a solution of the acetate described in Step 3 (50.4 g, 150 mmol) in 1.5 L of 3:1 THF/water at 0 °C was added lithium hydroxide monohydrate (18.9 g, 450 mmol). After one hour, the reaction was concentrated *in vacuo*, diluted with EtOAc (3 L), and washed with water, sat. aq. NaHCO<sub>3</sub> and brine. The solution was then dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated *in vacuo* to provide the crude product as a pale yellow fluffy solid which was sufficiently pure for use in the next step without further purification.

15 Step 5: Preparation of 5-(Chloromethyl)-1-(4-cyanobenzyl)-  
imidazole hydrochloride

The alcohol described in Step 4 (15.5 g, 72.9 mmol) was dissolved in thionyl chloride (150 mL) (exothermic!), and the reaction heated at 50°C under argon for 48h. Excess thionyl chloride was distilled *in vacuo*, and the resulting solid stirred in methylene chloride for 2h. The solid was filtered and dried *in vacuo*, giving the title compound.


Step 6: Preparation of 5-(5-(5-azabenzimidazolyl)methyl)-1-(4-cyanobenzyl)imidazole hydrochloride

To a solution of 5-azabenzimidazole (119 mg, 1.00 mmol) in 3 mL of dry DMF was added sodium hydride (88.0 mg, 2.20 mmol, 60% dispersion in mineral oil) at room temperature. After one hour, the solution was cooled to -50 °C and the chloride described in Step 5 (268 mg, 1.00 mmol) was added as a solid. The reaction was slowly warmed to room temperature over 16 hours, poured onto water, and extracted with methylene chloride (3 x 10 mL). The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated *in vacuo* to provide a yellow oil. The crude product was purified by column

chromatography (10-30% MeOH/CHCl<sub>3</sub>) and then treated with HCl to provide the title compound as a white solid. MS (FAB) m+1 = 315. elemental analysis for C<sub>18</sub>H<sub>14</sub>N<sub>6</sub> · 3.70 HCl · .75 H<sub>2</sub>O calc. C, 46.71; H, 4.18; N, 18.16; found C, 46.73; H, 4.17; N, 17.95.

5

#### EXAMPLE 5



#### 10 5-[4-(4-Azabenzimidazolyl)methyl]-1-(4-cyanobenzyl)imidazole hydrochloride

The title compound was prepared in a similar manner as described in Example 4, except 4-azabenzimidazole was used in Step 6. MS (FAB) m+1 = 315. elemental analysis for C<sub>18</sub>H<sub>14</sub>N<sub>6</sub> · 4.05 HCl · .45 EtOAc calc. C, 47.40; H, 4.35; N, 16.75; found C, 47.34; H, 4.34; N, 16.76.

15

#### EXAMPLE 6

##### In vitro inhibition of ras farnesyl transferase

20                   *Assays of farnesyl-protein transferase.* Partially purified bovine FPTase and Ras peptides (Ras-CVLS (SEQ.ID.NO.: 11), Ras-CVIM (SEQ.ID.NO.: 1) and Ras-CAIL (SEQ.ID.NO.: 12)) were prepared as described by Schaber *et al.*, *J. Biol. Chem.* 265:14701-14704 (1990), Pompliano, *et al.*, *Biochemistry* 31:3800 (1992) and  
25   Gibbs *et al.*, *PNAS* U.S.A. 86:6630-6634 (1989), respectively. Bovine FPTase was assayed in a volume of 100 ml containing 100 mM N-(2-hydroxy ethyl) piperazine-N'-(2-ethane sulfonic acid) (HEPES), pH 7.4, 5 mM MgCl<sub>2</sub>, 5 mM dithiothreitol (DTT), 100 mM [<sup>3</sup>H]-farnesyl diphosphate ([<sup>3</sup>H]-FPP; 740 CBq/mmol, New England Nuclear), 650 nM

Ras-CVLS and 10 mg/ml FPTase at 31°C for 60 min. Reactions were initiated with FPTase and stopped with 1 ml of 1.0 M HCL in ethanol. Precipitates were collected onto filter-mats using a TomTec Mach II cell harvester, washed with 100% ethanol, dried and counted in an LKB b-plate counter. The assay was linear with respect to both substrates, FPTase levels and time; less than 10% of the [<sup>3</sup>H]-FPP was utilized during the reaction period. Purified compounds were dissolved in 100% dimethyl sulfoxide (DMSO) and were diluted 20-fold into the assay. Percentage inhibition is measured by the amount of incorporation of radioactivity in the presence of the test compound when compared to the amount of incorporation in the absence of the test compound.

Human FPTase was prepared as described by Omer *et al.*, Biochemistry 32:5167-5176 (1993). Human FPTase activity was assayed as described above with the exception that 0.1% (w/v) polyethylene glycol 20,000, 10 mM ZnCl<sub>2</sub> and 100 nM Ras-CVIM were added to the reaction mixture. Reactions were performed for 30 min., stopped with 100 ml of 30% (v/v) trichloroacetic acid (TCA) in ethanol and processed as described above for the bovine enzyme.

The compounds of the instant invention described in the above Examples were tested for inhibitory activity against human FPTase by the assay described above and were found to have IC<sub>50</sub> of <50 μM.

25

### EXAMPLE 7

#### Modified *In vitro* GGTase inhibition assay

The modified geranylgeranyl-protein transferase inhibition assay is carried out at room temperature. A typical reaction contains (in a final volume of 50 mL): [<sup>3</sup>H]geranylgeranyl diphosphate, biotinylated Ras peptide, 50 mM HEPES, pH 7.5, a modulating anion (for example 10 mM glycerophosphate or 5mM ATP), 5 mM MgCl<sub>2</sub>, 10 mM ZnCl<sub>2</sub>, 0.1% PEG (15-20,000), 2 mM dithiothreitol, and geranylgeranyl-protein transferase type I (GGTase). The GGTase-type I enzyme



employed in the assay is prepared as described in U.S. Pat. No. 5,470,832, incorporated by reference. The Ras peptide is derived from the K4B-Ras protein and has the following sequence: biotinyl-GKKKKKKSKTKCVIM (single amino acid code) (SEQ.ID.NO.: 13).

- 5 Reactions are initiated by the addition of GGTase and stopped at timed intervals (typically 15 min) by the addition of 200  $\mu$ L of a 3 mg/mL suspension of streptavidin SPA beads (Scintillation Proximity Assay beads, Amersham) in 0.2 M sodium phosphate, pH 4, containing 50 mM EDTA, and 0.5% BSA. The quenched reactions are allowed to stand  
10 for 2 hours before analysis on a Packard TopCount scintillation counter.

- For inhibition studies, assays are run as described above, except inhibitors are prepared as concentrated solutions in 100% dimethyl sulfoxide and then diluted 25-fold into the enzyme assay mixture.  $IC_{50}$  values are determined with Ras peptide near  $K_M$   
15 concentrations. Enzyme and nonsaturating substrate conditions for inhibitor  $IC_{50}$  determinations are as follows: 75 pM GGTase-I, 1.6 mM Ras peptide, 100 nM geranylgeranyl diphosphate.

### EXAMPLE 8

20

#### Cell-based *in vitro* ras prenylation assay

- The cell lines used in this assay consist of either Rat1 or NIH3T3 cells transformed by either viral H-ras; an N-ras chimeric gene in which the C-terminal hypervariable region of viral-H-ras was  
25 substituted with the corresponding region from the N-ras gene; or ras-CVLL, a viral-H-ras mutant in which the C-terminal exon encodes leucine instead of serine, making the encoded protein a substrate for geranylgeranylation by GGTase-I. The assay can also be performed using cell lines transformed with human H-ras, N-ras or K4B-ras.  
30 The assay is performed essentially as described in DeClue, J.E. et al., Cancer Research 51:712-717, (1991). Cells in 10 cm dishes at 50-75% confluency are treated with the test compound(s) (final concentration of solvent, methanol or dimethyl sulfoxide, is 0.1%). After 4 hours at 37°C, the cells are labelled in 3 ml methionine-free DMEM supple-

mented with 10% regular DMEM, 2% fetal bovine serum, 400 mCi[<sup>35</sup>S]methionine (1000 Ci/mmol) and test compound(s). Cells treated with lovastatin, a compound that blocks Ras processing in cells by inhibiting the rate-limiting step in the isoprenoid biosynthetic pathway (Hancock, J.F. et al. *Cell*, 57:1167 (1989); DeClue, J.E. et al. *Cancer Res.*, 51:712 (1991); Sinensky, M. et al. *J. Biol. Chem.*, 265:19937 (1990)), serve as a positive control in this assay. After an additional 20 hours, the cells are lysed in 1 ml lysis buffer (1% NP40/20 mM HEPES, pH 7.5/5 mM MgCl<sub>2</sub>/1mM DTT/10 mg/ml aprotinen/2 mg/ml leupeptin/2 mg/ml antipain/0.5 mM PMSF) and the lysates cleared by centrifugation at 100,000 x g for 45 min. Alternatively, four hours after the additon of the labelling media, the media is removed, the cells washed, and 3 ml of media containing the same or a different test compound added. Following an additional 16 hour incubation, the lysis is carried out as above. Aliquots of lysates containing equal numbers of acid-precipitable counts are bought to 1 ml with IP buffer (lysis buffer lacking DTT) and immunoprecipitated with the ras-specific monoclonal antibody Y13-259 (Furth, M.E. et al., *J. Virol.* 43:294-304, (1982)). Following a 2 hour antibody incubation at 4°C, 200 ml of a 25% suspension of protein A-Sepharose coated with rabbit anti rat IgG is added for 45 min. The immunoprecipitates are washed four times with IP buffer (20 nM HEPES, pH 7.5/1 mM EDTA/1% Triton X-100.0.5% deoxycholate/0.1%/SDS/0.1 M NaCl) boiled in SDS-PAGE sample buffer and loaded on 13% acrylamide gels. When the dye front reached the bottom, the gel is fixed, soaked in Enlightening, dried and autoradiographed. The intensities of the bands corresponding to prenylated and nonprenylated Ras proteins are compared to determine the percent inhibition of prenyl transfer to protein.

30

### EXAMPLE 9

#### Cell-based *in vitro* anchorage independent growth assay (SALSA)

SALSA (Soft Agar-Like Surrogate Assay) measures the inhibition of anchorage-independent growth by prenyl-transferase inhibitors. Only transformed cells are able to grow anchorage-

35

independently in the SALSA format. Additionally, cells growing in the SALSA format grow in clumps, resembling the colonies formed in soft agar. SALSA may be used to measure the growth inhibition by prenyl-transferase inhibitors in a variety of

5 transformed cell lines, including Rat1 fibroblasts transformed with viral-H-ras (H-ras/rat1), as well as a panel of human tumor cell lines (HTL's).

SALSA is performed in 96-well plates that are coated with a thin film of the polymer, PolyHEMA (Poly(2-hydroxyethyl methacrylate)), which prevents cells from attaching to the plate. Rat1 fibroblast cells transformed with v-Ha-ras (this cell line has been deposited in the ATCC on August 19, 1997 under the terms of the Budapest convention and has been given a designation of ATCC CRL 12387) are seeded at 5000 cells/well, grown for 4 hr, then  
10 vehicle or half-log dilutions of test compound (in either an 8 or 12 point titration) are added. The cells are then grown for 6 days at 37 degrees, without changing the growth media or adding fresh compound. At day 6, cell growth is assessed via a colorimetric assay that measures the cleavage of the tetrazolium dye, MTT, to an  
15 insoluble purple formazan, a reaction dependent upon mitochondrial dehydrogenases. At day 6, the cells are incubated for 4 hr with 0.5 mg/ml MTT, and then SDS is added to 9% w/v to lyse the cells and solubilize the insoluble MTT-formazan. The amount of MTT metabolism is quantitated via spectrophotometric detection at 570  
20 nM. Dose-inhibition curves and  $IC_{50}$ 's are determined.

### EXAMPLE 10

#### Construction of SEAP reporter plasmid pDSE100

30 The SEAP reporter plasmid, pDSE100 was constructed by ligating a restriction fragment containing the SEAP coding sequence into the plasmid pCMV-RE-AKI. The SEAP gene is derived from the plasmid pSEAP2-Basic (Clontech, Palo Alto, CA). The plasmid pCMV-RE-AKI was constructed by Deborah Jones (Merck) and contains 5

sequential copies of the 'dyad symmetry response element' cloned upstream of a 'CAT-TATA' sequence derived from the cytomegalovirus immediate early promoter. The plasmid also contains a bovine growth hormone poly-A sequence.

5           The plasmid, pDSE100 was constructed as follows. A restriction fragment encoding the SEAP coding sequence was cut out of the plasmid pSEAP2-Basic using the restriction enzymes EcoR1 and HpaI. The ends of the linear DNA fragments were filled in with the Klenow fragment of E. coli DNA Polymerase I. The 'blunt ended' DNA  
10 containing the SEAP gene was isolated by electrophoresing the digest in an agarose gel and cutting out the 1694 base pair fragment. The vector plasmid pCMV-RE-AKI was linearized with the restriction enzyme Bgl-II and the ends filled in with Klenow DNA Polymerase I. The SEAP DNA fragment was blunt end ligated into the pCMV-RE-AKI vector  
15 and the ligation products were transformed into DH5-alpha E. coli cells (Gibco-BRL). Transformants were screened for the proper insert and then mapped for restriction fragment orientation. Properly oriented recombinant constructs were sequenced across the cloning junctions to verify the correct sequence. The resulting plasmid contains the SEAP  
20 coding sequence downstream of the DSE and CAT-TATA promoter elements and upstream of the BGH poly-A sequence.

#### Cloning of a Myristylated viral-H-ras expression plasmid

25   A DNA fragment containing viral-H-ras can be PCR'd from plasmid "H-1" (Ellis R. et al. J. Virol. 36, 408, 1980) using the following oligos.

Sense strand:

5'TCTCCTCGAGGCCACCATGGGGAGTAGCAAGAGCAAGCCTAA  
30 GGACCCAGCCAGCGCCGGATGACAGAATACAAGCTTGTGGTG  
G 3'. (SEQ.ID.NO.: 14)

Antisense: 5'CACATCTAGATCAGGACAGCACAGACTTGCAGC 3'.  
(SEQ.ID.NO.: 15)

A sequence encoding the first 15 aminoacids of the v-src gene, containing a myristylation site, is incorporated into the sense strand oligo. The sense strand oligo also optimizes the 'Kozak' translation initiation sequence immediately 5' to the ATG start site. To prevent prenylation at the viral-*ras* C-terminus, cysteine 186 would be mutated to a serine by substituting a G residue for a C residue in the C-terminal antisense oligo. The PCR primer oligos introduce an XhoI site at the 5' end and a XbaI site at the 3' end. The XhoI-XbaI fragment can be ligated into the mammalian expression plasmid pCI (Promega) cut with XhoI and XbaI. This results in a plasmid in which the recombinant myr-viral-H-*ras* gene is constitutively transcribed from the CMV promoter of the pCI vector.

15 Cloning of a viral-H-*ras*-CVLL expression plasmid

A viral-H-*ras* clone with a C-terminal sequence encoding the amino acids CVLL can be cloned from the plasmid "H-1" (Ellis R. et al. J. Virol. 36, 408, 1980) by PCR using the following oligos.

20

Sense strand:

5'TCTCCTCGAGGCCACCATGACAGAATACAAGCTTGTGGTGG-3' (SEQ.ID.NO.: 16)

25 Antisense strand:

5'CACTCTAGACTGGTGTTCAGAGCAGCACACACTTGCAGC-3' (SEQ.ID.NO.: 17)

30 The sense strand oligo optimizes the 'Kozak' sequence and adds an XhoI site. The antisense strand mutates serine 189 to leucine and adds an XbaI site. The PCR fragment can be trimmed with XhoI and XbaI and ligated into the XhoI-XbaI cut vector pCI (Promega). This results in a plasmid in which the mutated viral-H-*ras*-CVLL gene is constitutively transcribed from the CMV promoter of the pCI vector.

Cloning of c-H-ras-Leu61 expression plasmid

5 The human c-H-ras gene can be PCR'd from a human cerebral cortex cDNA library (Clontech) using the following oligonucleotide primers.

Sense strand:

5'-GAGAGAATTCGCCACCATGACGGAATATAAGCTGGTGG-3'  
(SEQ.ID.NO.: 18)

10

Antisense strand:

5'-GAGAGTCGACGCGTCAGGAGAGCACACACTTGC-3'  
(SEQ.ID.NO.: 19)

15 The primers will amplify a c-H-ras encoding DNA fragment with the primers contributing an optimized 'Kozak' translation start sequence, an EcoRI site at the N-terminus and a Sal I site at the C-terminal end. After trimming the ends of the PCR product with EcoRI and Sal I, the c-H-ras fragment can be ligated into an EcoRI -Sal I cut  
20 mutagenesis vector pAlter-1 (Promega). Mutation of glutamine-61 to a leucine can be accomplished using the manufacturer's protocols and the following oligonucleotide:

5'-CCGCCGGCCTGGAGGAGTACAG-3' (SEQ.ID.NO.: 20)

25

After selection and sequencing for the correct nucleotide substitution, the mutated c-H-ras-Leu61 can be excised from the pAlter-1 vector, using EcoRI and Sal I, and be directly ligated into the vector pCI (Promega) which has been digested with EcoRI and Sal I. The new  
30 recombinant plasmid will constitutively transcribe c-H-ras-Leu61 from the CMV promoter of the pCI vector.

Cloning of a c-N-ras-Val-12 expression plasmid

The human c-N-ras gene can be PCR'd from a human cerebral cortex cDNA library (Clontech) using the following oligonucleotide primers.

5

Sense strand:

5'-GAGAGAATTCGCCACCATGACTGAGTACAAACTGGTGG-3'  
(SEQ.ID.NO.: 21)

10 Antisense strand:

5'-GAGAGTCGACTTGTTACATCACCACACATGGC-3'  
(SEQ.ID.NO.: 22)

15 The primers will amplify a c-N-ras encoding DNA fragment with the primers contributing an optimized 'Kozak' translation start sequence, an EcoRI site at the N-terminus and a Sal I site at the C-terminal end.

After trimming the ends of the PCR product with EcoRI and Sal I, the c-N-ras fragment can be ligated into an EcoRI -Sal I cut mutagenesis vector pAlter-1 (Promega). Mutation of glycine-12 to a valine can be  
20 accomplished using the manufacturer's protocols and the following oligonucleotide:

5'-GTTGGAGCAGTTGGTGTGGG-3' (SEQ.ID.NO.: 23)

25 After selection and sequencing for the correct nucleotide substitution, the mutated c-N-ras-Val-12 can be excised from the pAlter-1 vector, using EcoRI and Sal I, and be directly ligated into the vector pCI (Promega) which has been digested with EcoRI and Sal I. The new recombinant plasmid will constitutively transcribe c-N-ras-Val-12 from  
30 the CMV promoter of the pCI vector.

### Cloning of a c-K-ras-Val-12 expression plasmid

The human c-K-ras gene can be PCR'd from a human cerebral cortex cDNA library (Clontech) using the following oligonucleotide primers.

5

Sense strand:

5'-GAGAGGTACCGCCACCATGACTGAATATAAACTTGTGG-3'  
(SEQ.ID.NO.: 24)

10 Antisense strand:

5'-CTCTGTCGACGTATTTACATAATTACACACTTTGTC-3'  
(SEQ.ID.NO.: 25)

15 The primers will amplify a c-K-ras encoding DNA fragment with the primers contributing an optimized 'Kozak' translation start sequence, a KpnI site at the N-terminus and a Sal I site at the C-terminal end. After trimming the ends of the PCR product with Kpn I and Sal I, the c-K-ras fragment can be ligated into a KpnI -Sal I cut mutagenesis vector pAlter-1 (Promega). Mutation of cysteine-12 to a valine can be  
20 accomplished using the manufacturer's protocols and the following oligonucleotide:

5'-GTAGTTGGAGCTGTTGGCGTAGGC-3' (SEQ.ID.NO.: 26)

25 After selection and sequencing for the correct nucleotide substitution, the mutated c-K-ras-Val-12 can be excised from the pAlter-1 vector, using KpnI and Sal I, and be directly ligated into the vector pCI (Promega) which has been digested with KpnI and Sal I. The new recombinant plasmid will constitutively transcribe c-K-ras-Val-12 from  
30 the CMV promoter of the pCI vector.

### SEAP assay

Human C33A cells (human epithelial carcinoma - ATTC collection) are seeded in 10cm tissue culture plates in DMEM + 10%



fetal calf serum + 1X Pen/Strep + 1X glutamine + 1X NEAA. Cells are grown at 37°C in a 5% CO<sub>2</sub> atmosphere until they reach 50 -80% of confluency.

The transient transfection is performed by the CaPO<sub>4</sub> method (Sambrook et al., 1989). Thus, expression plasmids for H-*ras*, N-*ras*, K-*ras*, Myr-*ras* or H-*ras*-CVLL are co-precipitated with the DSE-SEAP reporter construct. For 10cm plates 600ml of CaCl<sub>2</sub>-DNA solution is added dropwise while vortexing to 600ml of 2X HBS buffer to give 1.2ml of precipitate solution (see recipes below). This is allowed to sit at room temperature for 20 to 30 minutes. While the precipitate is forming, the media on the C33A cells is replaced with DMEM (minus phenol red; Gibco cat. # 31053-028)+ 0.5% charcoal stripped calf serum + 1X (Pen/Strep, Glutamine and nonessential aminoacids). The CaPO<sub>4</sub>-DNA precipitate is added dropwise to the cells and the plate rocked gently to distribute. DNA uptake is allowed to proceed for 5-6 hrs at 37°C under a 5% CO<sub>2</sub> atmosphere.

Following the DNA incubation period, the cells are washed with PBS and trypsinized with 1ml of 0.05% trypsin. The 1 ml of trypsinized cells is diluted into 10ml of phenol red free DMEM + 0.2% charcoal stripped calf serum + 1X (Pen/Strep, Glutamine and NEAA ). Transfected cells are plated in a 96 well microtiter plate (100ml/well) to which drug, diluted in media, has already been added in a volume of 100ml. The final volume per well is 200ml with each drug concentration repeated in triplicate over a range of half-log steps.

Incubation of cells and drugs is for 36 hrs at 37° under CO<sub>2</sub>. At the end of the incubation period, cells are examined microscopically for evidence of cell distress. Next, 100ml of media containing the secreted alkaline phosphatase is removed from each well and transferred to a microtube array for heat treatment at 65°C for 1 hr to inactivate endogenous alkaline phosphatases (but not the heat stable secreted phosphatase).

The heat treated media is assayed for alkaline phosphatase by a luminescence assay using the luminescence reagent CSPD®

- (Tropix, Bedford, Mass.). A volume of 50 ml media is combined with 200 ml of CSPD cocktail and incubated for 60 minutes at room temperature. Luminescence is monitored using an ML2200 microplate luminometer (Dynatech). Luminescence reflects the level of activation of the fos reporter construct stimulated by the transiently expressed protein.

DNA-CaPO<sub>4</sub> precipitate for 10cm. plate of cells

- |    |                                  |       |
|----|----------------------------------|-------|
| 10 | Ras expression plasmid (1mg/ml)  | 10ml  |
|    | DSE-SEAP Plasmid (1mg/ml)        | 2ml   |
|    | Sheared Calf Thymus DNA (1mg/ml) | 8ml   |
|    | 2M CaCl <sub>2</sub>             | 74ml  |
|    | dH <sub>2</sub> O                | 506ml |

15

2X HBS Buffer

- |    |  |
|----|--|
|    | 280mM NaCl   |
|    | 10mM KCl   |
| 20 | 1.5mM Na <sub>2</sub> HPO <sub>4</sub> 2H <sub>2</sub> O |
|    | 12mM dextrose  |
|    | 50mM HEPES   |
|    | Final pH = 7.05  |

- 25 Luminescence Buffer (26ml)

- |    |                           |       |
|----|---------------------------|-------|
|    | Assay Buffer              | 20ml  |
|    | Emerald Reagent™ (Tropix) | 2.5ml |
|    | 100mM homoarginine        | 2.5ml |
| 30 | CSPD Reagent® (Tropix)    | 1.0ml |

Assay Buffer

Add 0.05M Na<sub>2</sub>CO<sub>3</sub> to 0.05M NaHCO<sub>3</sub> to obtain pH 9.5. Make 1mM in MgCl<sub>2</sub>


## EXAMPLE 11

### *In vivo* tumor growth inhibition assay (nude mouse)

- 5                    *In vivo* efficacy as an inhibitor of the growth of cancer cells may be confirmed by several protocols well known in the art. Examples of such *in vivo* efficacy studies are described by N. E. Kohl et al. (*Nature Medicine*, 1:792-797 (1995)) and N. E. Kohl et al. (*Proc. Nat. Acad. Sci. U.S.A.*, 91:9141-9145 (1994)).
- 10                   Rodent fibroblasts transformed with oncogenically mutated human Ha-*ras* or Ki-*ras* ( $10^6$  cells/animal in 1 ml of DMEM salts) are injected subcutaneously into the left flank of 8-12 week old female nude mice (Harlan) on day 0. The mice in each oncogene group are randomly assigned to a vehicle, compound or combination treatment
- 15                   group. Animals are dosed subcutaneously starting on day 1 and daily for the duration of the experiment. Alternatively, the farnesyl-protein transferase inhibitor may be administered by a continuous infusion pump. Compound, compound combination or vehicle is delivered in a total volume of 0.1 ml. Tumors are excised and weighed when all of
- 20                   the vehicle-treated animals exhibited lesions of 0.5 - 1.0 cm in diameter, typically 11-15 days after the cells were injected. The average weight of the tumors in each treatment group for each cell line is calculated.

WHAT IS CLAIMED IS:

1. A compound which inhibits farnesyl-protein  
 5 transferase of the formula I:



wherein:

R1a, R1b and R1c are independently selected from:

- 10 a) hydrogen,  
 b) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, C3-C10 cycloalkyl, C2-C6 alkenyl, C2-C6 alkynyl, R8O-, R9S(O)m-, R8C(O)NR8-, CN, NO2, (R8)2N-C(NR8)-, R8C(O)-, R8OC(O)-, N3, -N(R8)2, or  
 15 R9OC(O)NR8-,  
 c) C1-C6 alkyl unsubstituted or substituted by unsubstituted or substituted aryl, heterocyclic, C3-C10 cycloalkyl, C2-C6 alkenyl, C2-C6 alkynyl, R8O-, R9S(O)m-, R8C(O)NR8-, CN, (R8)2N-C(NR8)-, R8C(O)-, R8OC(O)-, N3, -N(R8)2, or R9OC(O)-NR8-,  
 20 provided that R1a is not unsubstituted or substituted imidazolyl;

R2a, R2b' and R2b'' are independently hydrogen or  
 -(CR112)vA3(CR122)wR13; or

- 25 R2b' and R2b'' are combined as O;

R<sup>3a</sup> and R<sup>3b</sup> are independently selected from:

- a) hydrogen,
- b) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, unsubstituted or substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl, unsubstituted or substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, halogen, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>9</sup>C(O)O-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, NO<sub>2</sub>, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,
- c) unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl,
- d) substituted C<sub>1</sub>-C<sub>6</sub> alkyl wherein the substituent on the substituted C<sub>1</sub>-C<sub>6</sub> alkyl is selected from unsubstituted or substituted aryl, unsubstituted or substituted heterocyclic, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, and R<sup>9</sup>OC(O)-NR<sup>8</sup>-;

R<sup>4</sup> is independently selected from:

- a) hydrogen,
- b) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, perfluoroalkyl, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, R<sup>8</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-, and
- c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl, heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, perfluoroalkyl, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NH-, CN, H<sub>2</sub>N-C(NH)-, R<sup>8</sup>C(O)-, R<sup>8</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>8</sup>OC(O)NH-,

provided that R<sup>4</sup> is not unsubstituted or substituted imidazolyl;

R<sup>5</sup> is independently selected from:




- a) hydrogen,
- b) C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, perfluoroalkyl, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, (R<sup>8</sup>)<sub>2</sub>N-C-(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, R<sup>8</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-, and
- c) C<sub>1</sub>-C<sub>6</sub> alkyl, unsubstituted or substituted by perfluoroalkyl, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, R<sup>8</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-;

R<sup>8</sup> is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, 2,2,2-trifluoroethyl and aryl;

15 R<sup>9</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

R<sup>10</sup> is selected from: H; R<sup>8</sup>C(O)-; R<sup>9</sup>S(O)<sub>m</sub>-; unsubstituted or substituted C<sub>1</sub>-4 alkyl, unsubstituted or substituted C<sub>3</sub>-6 cycloalkyl, unsubstituted or substituted heterocycle, unsubstituted or substituted aryl, substituted aroyl, unsubstituted or substituted heteroaroyl, substituted arylsulfonyl, unsubstituted or substituted heteroarylsulfonyl, wherein the substituted group is substituted with one or two substituents selected from:

- a) C<sub>1</sub>-4 alkoxy,
- b) aryl or heterocycle,
- c) halogen,
- d) HO,



h)  $\text{N}(\text{R}^8)_2$ , or

i) C3-6 cycloalkyl;

5

$\text{R}^{11}$  and  $\text{R}^{12}$  are independently selected from:

- a) hydrogen,
- b) C1-C6 alkyl unsubstituted or substituted by C2-C20 alkenyl,  $\text{R}^8\text{O}-$ ,  $\text{R}^9\text{S}(\text{O})_m-$ ,  $\text{R}^8\text{C}(\text{O})\text{NR}^8-$ , CN,  $\text{N}_3$ ,  $(\text{R}^8)_2\text{N}-\text{C}(\text{NR}^8)-$ ,  $\text{R}^8\text{C}(\text{O})-$ ,  $-\text{N}(\text{R}^8)_2$ , or  $\text{R}^9\text{OC}(\text{O})\text{NR}^8-$ ,
- 10 c) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, C3-C10 cycloalkyl, C2-C20 alkenyl, halogen,  $\text{R}^8\text{O}-$ ,  $\text{R}^9\text{S}(\text{O})_m-$ ,  $\text{R}^8\text{C}(\text{O})\text{NR}^8-$ , CN,  $\text{NO}_2$ ,  $(\text{R}^8)_2\text{N}-\text{C}(\text{NR}^8)-$ ,
- 15 d) C1-C6 alkyl substituted with an unsubstituted or substituted group selected from aryl, heterocyclic and C3-C10 cycloalkyl;

20  $\text{R}^{13}$  is selected from:

- a) hydrogen,
- b) substituted or unsubstituted aryl, substituted or unsubstituted heterocycle, C3-C10 cycloalkyl, C2-C20 alkenyl, C2-C20 alkynyl, C1-C20 perfluoroalkyl, allyloxy,
- 25 F, Cl, Br,  $\text{R}^8\text{O}-$ ,  $\text{R}^9\text{S}(\text{O})_m-$ ,  $\text{R}^8\text{C}(\text{O})\text{NR}^8-$ , CN,  $\text{NO}_2$ ,  $\text{R}^8_2\text{N}-\text{C}(\text{NR}^8)-$ ,  $\text{R}^8\text{C}(\text{O})-$ ,  $\text{N}_3$ ,  $-\text{N}(\text{R}^8)_2$ ,  $(\text{R}^9)_2\text{NC}(\text{O})-$  or  $\text{R}^9\text{OC}(\text{O})\text{NR}^8-$ , and

- 5 c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by substituted or unsubstituted aryl, substituted or unsubstituted heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>2</sub>-C<sub>20</sub> alkynyl, C<sub>2</sub>-C<sub>20</sub> perfluoroalkyl, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NH-, CN, H<sub>2</sub>N-C(NH)-, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NH-;

10 A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR<sup>8</sup>-, -NR<sup>8</sup>C(O)-, O, -N(R<sup>8</sup>)-, -S(O)<sub>2</sub>N(R<sup>8</sup>)-, -N(R<sup>8</sup>)S(O)<sub>2</sub>-, or -S(O)<sub>m</sub>;

15 A<sup>3</sup> are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR<sup>10</sup>-, -NR<sup>10</sup>C(O)-, O, -N(R<sup>10</sup>)-, -S(O)<sub>2</sub>N(R<sup>10</sup>)-, -N(R<sup>10</sup>)S(O)<sub>2</sub>-, or -S(O)<sub>m</sub>;

20 J, K and L are independently selected from: S, N, O, NH and CH;

V is selected from:

- 25 a) hydrogen,  
b) heterocycle,  
c) aryl,  
d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N,  
and  
e) C<sub>2</sub>-C<sub>20</sub> alkenyl,

provided that V is not hydrogen if A<sup>1</sup> is -S(O)<sub>m</sub> and V is not hydrogen if A<sup>1</sup> is a bond, n is 0 and A<sup>2</sup> is -S(O)<sub>m</sub>;  
and provided that V is not imidazolyl;

30 W is a heterocycle;

X is a bond, -S(O)<sub>m</sub>-, O or -C(=O)-;

m is 0, 1 or 2;




- n is 0, 1, 2, 3 or 4;  
 p is 0, 1, 2, 3 or 4;  
 q is 0, 1, 2, 3 or 4, provided that q is not 0 or 1 if X is O;  
 r is 0 to 5, provided that r is 0 when V is hydrogen;  
 5 s is 1 or 2;  
 t is 0 or 1;  
 u is independently 0, 1 or 2;  
 v is 0, 1, 2, 3 or 4, provided that v is not 0 when A<sup>3</sup> is  
 -NR<sup>10</sup>C(O)-, O, -N(R<sup>10</sup>)-, -S(O)<sub>2</sub>N(R<sup>10</sup>)-,  
 10 -N(R<sup>10</sup>)S(O)<sub>2</sub>-, or S(O)<sub>m</sub>;  
 w is 0, 1, 2, 3 or 4; and

the dashed lines represent optional double bonds;

- 15 or an optical isomer or a pharmaceutically acceptable salt thereof.

2. The compound according to Claim 1 which inhibits farnesyl-protein transferase of the formula A:



20

wherein:

R<sup>1a</sup>, R<sup>1b</sup> and R<sup>1c</sup> are independently selected from:

- a) hydrogen,  
 25 b) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl,

- C2-C6 alkynyl, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, R<sup>8</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,
- 5 c) C1-C6 alkyl unsubstituted or substituted by unsubstituted or substituted aryl, heterocyclic, C3-C10 cycloalkyl, C2-C6 alkenyl, C2-C6 alkynyl, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, R<sup>8</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)-NR<sup>8</sup>-,
- provided that R<sup>1a</sup> is not unsubstituted or substituted imidazolyl;
- 10 R<sup>2a</sup>, R<sup>2b'</sup> and R<sup>2b''</sup> are independently hydrogen or -(CR<sup>11</sup>)<sub>2</sub>VA<sup>3</sup>(CR<sup>12</sup>)<sub>2</sub>WR<sup>13</sup>; or R<sup>2b'</sup> and R<sup>2b''</sup> are combined as O;
- 15 R<sup>3a</sup> and R<sup>3b</sup> are independently selected from:
- a) hydrogen,
- b) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, unsubstituted or substituted C3-C10 cycloalkyl, unsubstituted or substituted C2-C6 alkenyl, C2-C6 alkynyl, halogen, C1-C6 perfluoroalkyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>9</sup>C(O)O-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, NO<sub>2</sub>, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,
- 20 c) unsubstituted C1-C6 alkyl,
- d) substituted C1-C6 alkyl wherein the substituent on the substituted C1-C6 alkyl is selected from unsubstituted or substituted aryl, unsubstituted or substituted heterocyclic, C3-C10 cycloalkyl, C2-C6 alkenyl, C2-C6 alkynyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, and R<sup>9</sup>OC(O)-NR<sup>8</sup>-;
- 25
- 30

R<sup>4</sup> is independently selected from:

- a) hydrogen,
  - b) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, perfluoroalkyl, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, R<sup>8</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-, and
  - c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl, heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, perfluoroalkyl, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NH-, CN, H<sub>2</sub>N-C(NH)-, R<sup>8</sup>C(O)-, R<sup>8</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>8</sup>OC(O)NH-,
- provided that R<sup>4</sup> is not unsubstituted or substituted imidazolyl;

R<sup>5</sup> is independently selected from:

- a) hydrogen,
- b) C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, perfluoroalkyl, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, (R<sup>8</sup>)<sub>2</sub>N-C-(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, R<sup>8</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-, and
- c) C<sub>1</sub>-C<sub>6</sub> alkyl, unsubstituted or substituted by perfluoroalkyl, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, R<sup>8</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-;




R<sup>8</sup> is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, 2,2,2-trifluoroethyl and aryl;

R<sup>9</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

R<sup>10</sup> is selected from: H; R<sup>8</sup>C(O)-; R<sup>9</sup>S(O)<sub>m</sub>-; unsubstituted or substituted C<sub>1</sub>-4 alkyl, unsubstituted or substituted C<sub>3</sub>-6 cycloalkyl, unsubstituted or substituted heterocycle, unsubstituted or substituted

aryl, substituted aroyl, unsubstituted or substituted heteroaroyl, substituted arylsulfonyl, unsubstituted or substituted heteroarylsulfonyl, wherein the substituted group is substituted with one or two substituents selected from:

- 5           a) C<sub>1</sub>-4 alkoxy,  
              b) aryl or heterocycle,  
              c) halogen,  
              d) HO,



10

- h) N(R<sup>8</sup>)<sub>2</sub>, or  
              i) C<sub>3</sub>-6 cycloalkyl;

R<sup>11</sup> and R<sup>12</sup> are independently selected from:

- 15           a) hydrogen,  
              b) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by C<sub>2</sub>-C<sub>20</sub> alkenyl,  
                   R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, N<sub>3</sub>,  
                   (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,  
              c) unsubstituted or substituted aryl, unsubstituted or  
 20           substituted heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-  
                   C<sub>20</sub> alkenyl, halogen, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-,  
                   R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-,  
                   R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-, and  
              d) C<sub>1</sub>-C<sub>6</sub> alkyl substituted with an unsubstituted or  
 25           substituted group selected from aryl, heterocyclic and  
                   C<sub>3</sub>-C<sub>10</sub> cycloalkyl;

R<sup>13</sup> is selected from:

- a) hydrogen,
- b) substituted or unsubstituted aryl, substituted or  
 5 unsubstituted heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>20</sub>  
 alkenyl, C<sub>2</sub>-C<sub>20</sub> alkynyl, C<sub>1</sub>-C<sub>20</sub> perfluoroalkyl, allyloxy,  
 F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>,  
 R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, (R<sup>9</sup>)<sub>2</sub>NC(O)- or  
 R<sup>9</sup>OC(O)NR<sup>8</sup>-, and
- c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by substituted or  
 10 unsubstituted aryl, substituted or unsubstituted heterocycle,  
 C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>2</sub>-C<sub>20</sub> alkynyl, C<sub>2</sub>-  
 C<sub>20</sub> perfluoroalkyl, F, Cl, Br, R<sup>8</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-,  
 R<sup>8</sup>C(O)NH-, CN, H<sub>2</sub>N-C(NH)-, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or  
 15 R<sup>9</sup>OC(O)NH-;

A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond, -CH=CH-, -C≡C-,  
 -C(O)-, -C(O)NR<sup>8</sup>-, -NR<sup>8</sup>C(O)-, O, -N(R<sup>8</sup>)-,  
 -S(O)<sub>2</sub>N(R<sup>8</sup>)-, -N(R<sup>8</sup>)S(O)<sub>2</sub>-, or -S(O)<sub>m</sub>;

20

A<sup>3</sup> are independently selected from: a bond, -CH=CH-,  
 -C≡C-, -C(O)-, -C(O)NR<sup>10</sup>-, -NR<sup>10</sup>C(O)-, O, -N(R<sup>10</sup>)-,  
 -S(O)<sub>2</sub>N(R<sup>10</sup>)-, -N(R<sup>10</sup>)S(O)<sub>2</sub>-, or -S(O)<sub>m</sub>;

25 J, K and L are independently selected from: S, N, O, NH and CH;

V is selected from:

- a) hydrogen,
- b) heterocycle,
- c) aryl,
- d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are  
 30 replaced with a heteroatom selected from O, S, and N,  
 and
- e) C<sub>2</sub>-C<sub>20</sub> alkenyl,

provided that V is not hydrogen if  $A^1$  is  $-S(O)_m$  and V is not hydrogen if  $A^1$  is a bond, n is 0 and  $A^2$  is  $-S(O)_m$ ; and provided that V is not imidazolyl;

5 W is a heterocycle;

X is a bond,  $-S(O)_m$ -, O or  $-C(=O)-$ ;


m is 0, 1 or 2;  
 10 n is 0, 1, 2, 3 or 4;  
 p is 0, 1, 2, 3 or 4;  
 q is 0, 1, 2, 3 or 4, provided that q is not 0 or 1 if X is O;  
 r is 0 to 5, provided that r is 0 when V is hydrogen;  
 s is 1 or 2;  
 15 t is 0 or 1;  
 u is independently 0, 1 or 2;  
 v is 0, 1, 2, 3 or 4, provided that v is not 0 when  $A^3$  is  $-NR^{10}C(O)-$ , O,  $-N(R^{10})-$ ,  $-S(O)_2N(R^{10})-$ ,  $-N(R^{10})S(O)_2-$ , or  $S(O)_m$ ;  
 20 w is 0, 1, 2, 3 or 4; and

the dashed lines represent optional double bonds;

or an optical isomer or a pharmaceutically acceptable salt thereof.

25

3. The compound according to Claim 2 which inhibits farnesyl-protein transferase of the formula A:



wherein:


R<sup>1a</sup> and R<sup>1c</sup> are independently selected from: hydrogen, C<sub>3</sub>-C<sub>10</sub>  
 5 cycloalkyl, R<sup>8</sup>O-, -N(R<sup>8</sup>)<sub>2</sub>, F or C<sub>1</sub>-C<sub>6</sub> alkyl;

R<sup>1b</sup> is independently selected from:

- a) hydrogen,
- b) unsubstituted or substituted aryl, unsubstituted or  
 10 substituted heterocycle, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, R<sup>8</sup>O-, -N(R<sup>8</sup>)<sub>2</sub>  
 or C<sub>2</sub>-C<sub>6</sub> alkenyl,
- c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by unsubstituted or  
 substituted aryl, heterocycle, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub>  
 alkenyl, R<sup>8</sup>O-, or -N(R<sup>8</sup>)<sub>2</sub>;

15

R<sup>2a</sup> and R<sup>2b'</sup> are independently selected from selected from:  
 H; C<sub>1</sub>-C<sub>6</sub> alkyl,



20 R<sup>3a</sup> and R<sup>3b</sup> are independently selected from:

- a) hydrogen,
- b) unsubstituted or substituted aryl, unsubstituted or  
 substituted heterocycle, unsubstituted or substituted  
 C<sub>3</sub>-C<sub>10</sub> cycloalkyl, unsubstituted or substituted C<sub>2</sub>-C<sub>6</sub>  
 25 alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, halogen, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,

- $R^9O-$ ,  $R^9S(O)_m-$ ,  $R^8C(O)NR^8-$ ,  $(R^8)_2NC(O)-$ ,  $R^9C(O)O-$ ,  
 $R^8_2N-C(NR^8)-$ ,  $CN$ ,  $NO_2$ ,  $R^8C(O)-$ ,  $N_3$ ,  $-N(R^8)_2$ , or  
 $R^9OC(O)NR^8-$ ,  
 5      c)    unsubstituted  $C_1-C_6$  alkyl,  
          d)    substituted  $C_1-C_6$  alkyl wherein the substituent on the  
               substituted  $C_1-C_6$  alkyl is selected from unsubstituted or  
               substituted aryl, unsubstituted or substituted heterocyclic,  
                $C_3-C_{10}$  cycloalkyl,  $C_2-C_6$  alkenyl,  $C_2-C_6$  alkynyl,  $R^9O-$ ,  
                $R^9S(O)_m-$ ,  $R^8C(O)NR^8-$ ,  $(R^8)_2NC(O)-$ ,  $R^8_2N-C(NR^8)-$ ,  
 10            $CN$ ,  $R^8C(O)-$ ,  $N_3$ ,  $-N(R^8)_2$ , and  $R^9OC(O)-NR^8-$ ;

$R^4$  is independently selected from:

- a)    hydrogen,  
 b)     $C_1-C_6$  alkyl,  $C_2-C_6$  alkenyl,  $C_2-C_6$  alkynyl,  $C_1-C_6$   
 15       perfluoroalkyl,  $F$ ,  $Cl$ ,  $R^8O-$ ,  $R^8C(O)NR^8-$ ,  $CN$ ,  $NO_2$ ,  
           $(R^8)_2N-C(NR^8)-$ ,  $R^8C(O)-$ ,  $R^8OC(O)-$ ,  $-N(R^8)_2$ , or  
           $R^9OC(O)NR^8-$ , and  
 c)     $C_1-C_6$  alkyl substituted by  $C_1-C_6$  perfluoroalkyl,  $R^8O-$ ,  
           $R^8C(O)NR^8-$ ,  $(R^8)_2N-C(NR^8)-$ ,  $R^8C(O)-$ ,  $R^8OC(O)-$ ,  
 20        $-N(R^8)_2$ , or  $R^9OC(O)NR^8-$ ;

$R^5$  is selected from:

- a)    hydrogen,  
 b)     $C_2-C_6$  alkenyl,  $C_2-C_6$  alkynyl,  $C_3-C_6$  cycloalkyl,  $C_1-C_6$   
 25       perfluoroalkyl,  $F$ ,  $Cl$ ,  $R^8O-$ ,  $R^9S(O)_m-$ ,  $R^8C(O)NR^8-$ ,  $CN$ ,  
           $NO_2$ ,  $(R^8)_2N-C(NR^8)-$ ,  $R^8C(O)-$ ,  $R^8OC(O)-$ ,  $-N(R^8)_2$ , or  
           $R^9OC(O)NR^8-$ , and  
 c)     $C_1-C_6$  alkyl unsubstituted or substituted by  $C_1-C_6$   
          perfluoroalkyl,  $F$ ,  $Cl$ ,  $R^8O-$ ,  $R^9S(O)_m-$ ,  $R^8C(O)NR^8-$ ,  $CN$ ,  
 30        $(R^8)_2N-C(NR^8)-$ ,  $R^8C(O)-$ ,  $R^8OC(O)-$ ,  $-N(R^8)_2$ , or  
           $R^9OC(O)NR^8-$ ;



$R^6$  and  $R^7$  are independently selected from:

H; C<sub>1</sub>-4 alkyl, C<sub>3</sub>-6 cycloalkyl, aryl, heterocycle,  
unsubstituted or substituted with one or two:



- a) C<sub>1</sub>-4 alkoxy,
- 5      b) halogen, or
- c) substituted or unsubstituted aryl or substituted  
or unsubstituted heterocycle,

10       $R^8$  is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, 2,2,2-trifluoroethyl and aryl;

$R^9$  is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

15       $R^{10}$  is selected from: H;  $R^8C(O)-$ ;  $R^9S(O)_m-$ ; unsubstituted or substituted C<sub>1</sub>-4 alkyl, unsubstituted or substituted C<sub>3</sub>-6 cycloalkyl, unsubstituted or substituted heterocycle, unsubstituted or substituted aryl, substituted aroyl, unsubstituted or substituted heteroaroyl, substituted arylsulfonyl, unsubstituted or substituted heteroarylsulfonyl, wherein the substituted group is substituted with one or two substituents  
20      selected from:

- a) C<sub>1</sub>-4 alkoxy,
- b) aryl or heterocycle,
- c) halogen,
- d) HO,



- h)  $N(R^8)_2$ , or

i) C<sub>3-6</sub> cycloalkyl;

A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond, -CH=CH-, -C≡C-,  
-C(O)-, -C(O)NR<sup>8</sup>-, O, -N(R<sup>8</sup>)-, or -S(O)<sub>m</sub>;

5

J, K and L are independently selected from: S, N, O, NH and CH;

V is selected from:

- 10           a) heterocycle selected from pyrrolidinyl, imidazolyl,  
pyridinyl, thiazolyl, indolyl, quinolinyl, isoquinolinyl, and  
thienyl, and  
             b) aryl;


15           W is a heterocycle selected from pyrrolidinyl, triazolyl, imidazolyl,  
pyridinyl, thiazolyl, indolyl, quinolinyl, or isoquinolinyl;

X is a bond, -S(O)<sub>m</sub>-, O or -C(=O)-;

- 20           m is           0, 1 or 2;  
n is           0, 1, 2, 3 or 4;  
p is           1, 2 or 3;  
q is           0, 1 or 2, provided that q is not 0 or 1 if X is O;  
r is           0 to 5, provided that r is 0 when V is hydrogen;  
s is           1 or 2;  
25           t is           1; and  
u is independently 0 or 1;

or an optical isomer or a pharmaceutically acceptable salt thereof.

30           4. The compound according to Claim 2 which inhibits  
farnesyl-protein transferase of the formula B:



wherein:


R<sup>1a</sup> and R<sup>1c</sup> are independently selected from: hydrogen, C<sub>3</sub>-C<sub>10</sub>  
 5 cycloalkyl, R<sup>8</sup>O-, -N(R<sup>8</sup>)<sub>2</sub>, F or C<sub>1</sub>-C<sub>6</sub> alkyl;

R<sup>1b</sup> is independently selected from:

- a) hydrogen,
- b) aryl, heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, R<sup>8</sup>O-, -N(R<sup>8</sup>)<sub>2</sub>, F or  
 10 C<sub>2</sub>-C<sub>6</sub> alkenyl,
- c) unsubstituted or substituted C<sub>1</sub>-C<sub>6</sub> alkyl wherein the  
 substituent on the substituted C<sub>1</sub>-C<sub>6</sub> alkyl is selected from  
 unsubstituted or substituted aryl, heterocycle, C<sub>3</sub>-C<sub>10</sub>  
 cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, R<sup>8</sup>O- and -N(R<sup>8</sup>)<sub>2</sub>;

15

R<sup>2a</sup> and R<sup>2b'</sup> are independently selected from selected from:  
 H; C<sub>1</sub>-C<sub>6</sub> alkyl,



20 R<sup>3a</sup> and R<sup>3b</sup> are independently selected from:

- a) hydrogen,
- b) unsubstituted or substituted aryl, unsubstituted or  
 substituted heterocycle, unsubstituted or substituted

- 5 C<sub>3</sub>-C<sub>10</sub> cycloalkyl, unsubstituted or substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, halogen, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>9</sup>C(O)O-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, NO<sub>2</sub>, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,
- 10 c) unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl,  
 d) substituted C<sub>1</sub>-C<sub>6</sub> alkyl wherein the substituent on the substituted C<sub>1</sub>-C<sub>6</sub> alkyl is selected from unsubstituted or substituted aryl, unsubstituted or substituted heterocyclic, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, and R<sup>9</sup>OC(O)-NR<sup>8</sup>-;

R<sup>4</sup> is independently selected from:

- 15 a) hydrogen,  
 b) aryl, substituted aryl, heterocycle, substituted heterocycle, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>8</sup>O-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,  
 20 and  
 c) C<sub>1</sub>-C<sub>6</sub> alkyl substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>8</sup>O-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-;
- 25 R<sup>5a</sup> and R<sup>5b</sup> are independently hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, cyclopropyl, trifluoromethyl and halogen;

R<sup>6</sup> and R<sup>7</sup> are independently selected from:

- 30 H; C<sub>1</sub>-4 alkyl, C<sub>3</sub>-6 cycloalkyl, aryl, heterocycle, unsubstituted or substituted with one or two:  
 a) C<sub>1</sub>-4 alkoxy,  
 b) halogen, or

- c) substituted or unsubstituted aryl or substituted or unsubstituted heterocycle;

5  $R^8$  is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, 2,2,2-trifluoroethyl, benzyl and aryl;

$R^9$  is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

10  $A^1$  and  $A^2$  are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR<sup>8</sup>-, O, -N(R<sup>8</sup>)-, or -S(O)<sub>m</sub>;

J, K and L are independently selected from: S, N, O, NH and CH;

V is selected from:

- 15 a) hydrogen,  
 b) heterocycle selected from pyrrolidinyl, imidazoliny, pyridinyl, thiazolyl, oxazolyl, indolyl, quinoliny, isoquinoliny, triazolyl and thienyl,  
 c) aryl,  
 20 d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and  
 e) C<sub>2</sub>-C<sub>20</sub> alkenyl, and


25 provided that V is not hydrogen if  $A^1$  is -S(O)<sub>m</sub> and V is not hydrogen if  $A^1$  is a bond, n is 0 and  $A^2$  is -S(O)<sub>m</sub>;

X is a bond, -S(O)<sub>m</sub>-, O or -C(=O)-;

30 m is 0, 1 or 2;  
 n is 0, 1, 2, 3 or 4;  
 p is 0, 1, 2, 3 or 4;  
 q is 0, 1 or 2, provided that q is not 0 or 1 if X is O;  
 r is 0 to 5, provided that r is 0 when V is hydrogen; and  
 u is independently 0 or 1;


or an optical isomer or pharmaceutically acceptable salt thereof.

5. The compound according to Claim 2 which inhibits  
 5 farnesyl-protein transferase of the formula C:



wherein:

- 10 R<sup>1a</sup> and R<sup>1c</sup> are independently selected from: hydrogen, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, R<sup>8</sup>O-, -N(R<sup>8</sup>)<sub>2</sub>, F or C<sub>1</sub>-C<sub>6</sub> alkyl;
- R<sup>1b</sup> is independently selected from:
- 15 a) hydrogen,
- b) aryl, heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, R<sup>8</sup>O-, -N(R<sup>8</sup>)<sub>2</sub>, F or C<sub>2</sub>-C<sub>6</sub> alkenyl,
- c) unsubstituted or substituted C<sub>1</sub>-C<sub>6</sub> alkyl wherein the substituent on the substituted C<sub>1</sub>-C<sub>6</sub> alkyl is selected from unsubstituted or substituted aryl, heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, R<sup>8</sup>O- and -N(R<sup>8</sup>)<sub>2</sub>;
- 20 R<sup>2a</sup> and R<sup>2b'</sup> are independently selected from selected from: H; C<sub>1</sub>-C<sub>6</sub> alkyl,



R<sup>3a</sup> and R<sup>3b</sup> are independently selected from:

- a) hydrogen,
- 5 b) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, unsubstituted or substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl, unsubstituted or substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, halogen, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>9</sup>C(O)O-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, NO<sub>2</sub>, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,
- 10 c) unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl,
- d) substituted C<sub>1</sub>-C<sub>6</sub> alkyl wherein the substituent on the substituted C<sub>1</sub>-C<sub>6</sub> alkyl is selected from unsubstituted or substituted aryl, unsubstituted or substituted heterocyclic, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, and R<sup>9</sup>OC(O)-NR<sup>8</sup>-;
- 15

20 R<sup>4</sup> is independently selected from:

- a) hydrogen,
- b) aryl, substituted aryl, heterocycle, substituted heterocycle, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>8</sup>O-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,
- 25 and
- c) C<sub>1</sub>-C<sub>6</sub> alkyl substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>8</sup>O-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-;
- 30

R<sup>5a</sup> and R<sup>5b</sup> are independently hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, cyclopropyl, trifluoromethyl and halogen;

$R^6$  and  $R^7$  are independently selected from:

H; C<sub>1-4</sub> alkyl, C<sub>3-6</sub> cycloalkyl, aryl, heterocycle,  
unsubstituted or substituted with one or two:

- 5      a) C<sub>1-4</sub> alkoxy,
- b) halogen, or
- c) substituted or unsubstituted aryl or substituted  
or unsubstituted heterocycle;

- 10     $R^8$  is independently selected from hydrogen, C<sub>1-C6</sub> alkyl, 2,2,2-trifluoroethyl, benzyl and aryl;

$R^9$  is independently selected from C<sub>1-C6</sub> alkyl and aryl;

- 15     $A^1$  and  $A^2$  are independently selected from: a bond, -CH=CH-, -C≡C-,  
-C(O)-, -C(O)NR<sup>8</sup>-, O, -N(R<sup>8</sup>)-, or -S(O)<sub>m</sub>;

J, K and L are independently selected from: S, N, O, NH and CH;

- 20    V is selected from:

- a) hydrogen,
- b) heterocycle selected from pyrrolidinyl, imidazoliny, pyridinyl, thiazolyl, oxazolyl, indolyl, quinoliny, isoquinoliny, triazolyl and thienyl,
- 25    c) aryl,
- d) C<sub>1-C20</sub> alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
- e) C<sub>2-C20</sub> alkenyl, and
- 30    provided that V is not hydrogen if  $A^1$  is -S(O)<sub>m</sub> and V is not hydrogen if  $A^1$  is a bond, n is 0 and  $A^2$  is -S(O)<sub>m</sub>;

X is a bond, -S(O)<sub>m</sub>-, O or -C(=O)-;




- m is 0, 1 or 2;  
 n is 0, 1, 2, 3 or 4;  
 p is 0, 1, 2, 3 or 4, provided that p is not 0 if X is a bond or O;  
 5 q is 0, 1 or 2, provided that q is not 0 or 1 if X is O;  
 r is 0 to 5, provided that r is 0 when V is hydrogen; and  
 u is independently 0 or 1;

or an optical isomer or pharmaceutically acceptable salt thereof.

10

6. The compound according to Claim 4 which inhibits farnesyl-protein transferase of the formula D:




15 wherein:

R<sup>1a</sup> and R<sup>1c</sup> are independently selected from: hydrogen, C<sub>3</sub>-C<sub>10</sub> cycloalkyl or C<sub>1</sub>-C<sub>6</sub> alkyl;


20 R<sup>1b</sup> is independently selected from:

- a) hydrogen,
- b) aryl, heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, R<sup>8</sup>O-, -N(R<sup>8</sup>)<sub>2</sub>, F or C<sub>2</sub>-C<sub>6</sub> alkenyl,
- c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl,
- 25 heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, R<sup>8</sup>O-, or



R<sup>2a</sup> and R<sup>2b'</sup> are independently selected from selected from:

5 H; C<sub>1</sub>-C<sub>6</sub> alkyl,



R<sup>3a</sup> and R<sup>3b</sup> are independently selected from:

- 10 a) hydrogen,  
b) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, unsubstituted or substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl, unsubstituted or substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, halogen, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>9</sup>C(O)O-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-,  
15 CN, NO<sub>2</sub>, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,  
c) unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl,  
d) substituted C<sub>1</sub>-C<sub>6</sub> alkyl wherein the substituent on the substituted C<sub>1</sub>-C<sub>6</sub> alkyl is selected from unsubstituted or substituted aryl, unsubstituted or substituted heterocyclic, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-,  
20 CN, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, and R<sup>9</sup>OC(O)-NR<sup>8</sup>-;

25 R<sup>4</sup> is independently selected from:

- a) hydrogen,  
b) aryl, substituted aryl, heterocycle, substituted heterocycle, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>8</sup>O-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>,  
30 (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,  
and

- c) C<sub>1</sub>-C<sub>6</sub> alkyl substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>8</sup>O-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-;

5 R<sup>5a</sup> and R<sup>5b</sup> are independently hydrogen, ethyl, cyclopropyl or methyl;

R<sup>6</sup> and R<sup>7</sup> are independently selected from:

H; C<sub>1</sub>-4 alkyl, C<sub>3</sub>-6 cycloalkyl, aryl, heterocycle,  
unsubstituted or substituted with one or two:

- 10 a) C<sub>1</sub>-4 alkoxy,  
b) halogen, or  
c) substituted or unsubstituted aryl or substituted  
or unsubstituted heterocycle;

15 R<sup>8</sup> is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, 2,2,2-trifluoroethyl, benzyl and aryl;

R<sup>9</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

20 A<sup>1</sup> is selected from: a bond, -C(O)-, O, -N(R<sup>8</sup>)-, or -S(O)<sub>m</sub>;

J, K and L are independently selected from: S, N, O, NH and CH;

X is a bond, -S(O)<sub>m</sub>-, O or -C(=O)-;

25

n is 0, 1 or 2; provided that n is not 0 or 1 if A<sup>1</sup> is a bond,  
O, -N(R<sup>8</sup>)-, or S(O)<sub>m</sub>;

m is 0, 1 or 2;

p is 0, 1, 2, 3 or 4;


30 q is 0, 1 or 2, provided that q is not 0 or 1 if X is O;

r is 1 or 2; and

u is independently 0 or 1;

or an optical isomer or pharmaceutically acceptable salt thereof.

7. The compound according to Claim 5 which inhibits farnesyl-protein transferase of the formula E:



5

wherein:


R<sup>1a</sup> and R<sup>1c</sup> are independently selected from: hydrogen, R<sup>8</sup>O-,  
-N(R<sup>8</sup>)<sub>2</sub>, F, C<sub>3</sub>-C<sub>10</sub> cycloalkyl or C<sub>1</sub>-C<sub>6</sub> alkyl;

10

R<sup>1b</sup> is independently selected from:

- a) hydrogen,
- b) aryl, heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, R<sup>8</sup>O-, -N(R<sup>8</sup>)<sub>2</sub>, F or C<sub>2</sub>-C<sub>6</sub> alkenyl,
- 15 c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl, heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, R<sup>8</sup>O-, or -N(R<sup>8</sup>)<sub>2</sub>;

20 R<sup>2a</sup> and R<sup>2b'</sup> are independently selected from selected from:  
H; C<sub>1</sub>-C<sub>6</sub> alkyl,



R<sup>3a</sup> and R<sup>3b</sup> are independently selected from:

- a) hydrogen,

- 5                   b)    unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, unsubstituted or substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl, unsubstituted or substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, halogen, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>9</sup>C(O)O-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, NO<sub>2</sub>, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,
- 10                   c)    unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl,
- d)    substituted C<sub>1</sub>-C<sub>6</sub> alkyl wherein the substituent on the substituted C<sub>1</sub>-C<sub>6</sub> alkyl is selected from unsubstituted or substituted aryl, unsubstituted or substituted heterocyclic, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, and R<sup>9</sup>OC(O)-NR<sup>8</sup>-;

15

R<sup>4</sup> is independently selected from:

- a)    hydrogen,
- b)    aryl, substituted aryl, heterocycle, substituted heterocycle, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>8</sup>O-, R<sup>8</sup>C(O)NR<sup>8</sup>-, CN, NO<sub>2</sub>, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,
- 20                   and
- c)    C<sub>1</sub>-C<sub>6</sub> alkyl substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>8</sup>O-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>N-C(NR<sup>8</sup>)-, R<sup>8</sup>C(O)-, -N(R<sup>8</sup>)<sub>2</sub>, or
- 25                   R<sup>9</sup>OC(O)NR<sup>8</sup>-;

R<sup>5a</sup> and R<sup>5b</sup> are independently hydrogen, ethyl, cyclopropyl or methyl;

- 30                   R<sup>6</sup> and R<sup>7</sup> are independently selected from:  
                   H; C<sub>1</sub>-4 alkyl, C<sub>3</sub>-6 cycloalkyl, aryl, heterocycle, unsubstituted or substituted with one or two:
- a)    C<sub>1</sub>-4 alkoxy,
- b)    halogen, or

c) substituted or unsubstituted aryl or substituted or unsubstituted heterocycle;

5  $R^8$  is independently selected from hydrogen,  $C_1$ - $C_6$  alkyl, 2,2,2-trifluoroethyl, benzyl and aryl;

$R^9$  is independently selected from  $C_1$ - $C_6$  alkyl and aryl;

10 J, K and L are independently selected from: S, N, O, NH and CH;

X is a bond,  $-S(O)_m-$ , O or  $-C(=O)-$ ;

n is 0 or 1;

m is 0, 1 or 2;

15 p is 0, 1, 2, 3 or 4, provided that p is not 0 if X is a bond or O;


q is 0, 1 or 2, provided that q is not 0 or 1 if X is O;

r is 1 or 2; and

u is independently 0 or 1;

20 or an optical isomer or pharmaceutically acceptable salt thereof.

8. The compound according to Claim 6 which inhibits farnesyl-protein transferase of the formula F:



25

wherein:


R<sup>1a</sup> and R<sup>1c</sup> are independently selected from: hydrogen, C<sub>3</sub>-C<sub>10</sub> cycloalkyl or C<sub>1</sub>-C<sub>6</sub> alkyl;

5

R<sup>1b</sup> is independently selected from:

- a) hydrogen,
  - b) aryl, heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, R<sup>8</sup>O-, -N(R<sup>8</sup>)<sub>2</sub> or F,
  - c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl,
- 10 heterocycle, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, R<sup>8</sup>O-, or -N(R<sup>8</sup>)<sub>2</sub>;

R<sup>2a</sup> and R<sup>2b'</sup> are independently selected from selected from: H; C<sub>1</sub>-C<sub>6</sub> alkyl,



15

R<sup>3a</sup> and R<sup>3b</sup> are independently selected from:

- a) hydrogen,
- b) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, unsubstituted or substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl, unsubstituted or substituted C<sub>2</sub>-C<sub>6</sub> alkenyl,
- 20 C<sub>2</sub>-C<sub>6</sub> alkynyl, halogen, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>9</sup>O-, R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>9</sup>C(O)O-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, NO<sub>2</sub>, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, or R<sup>9</sup>OC(O)NR<sup>8</sup>-,
- 25 c) unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl,
- d) substituted C<sub>1</sub>-C<sub>6</sub> alkyl wherein the substituent on the substituted C<sub>1</sub>-C<sub>6</sub> alkyl is selected from unsubstituted or substituted aryl, unsubstituted or substituted heterocyclic, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, R<sup>9</sup>O-,
- 30 R<sup>9</sup>S(O)<sub>m</sub>-, R<sup>8</sup>C(O)NR<sup>8</sup>-, (R<sup>8</sup>)<sub>2</sub>NC(O)-, R<sup>8</sup><sub>2</sub>N-C(NR<sup>8</sup>)-, CN, R<sup>8</sup>C(O)-, N<sub>3</sub>, -N(R<sup>8</sup>)<sub>2</sub>, and R<sup>9</sup>OC(O)-NR<sup>8</sup>-;

R<sup>5a</sup> and R<sup>5b</sup> are independently hydrogen, ethyl, cyclopropyl or methyl;

R<sup>6</sup> and R<sup>7</sup> are independently selected from:

H; C<sub>1-4</sub> alkyl, C<sub>3-6</sub> cycloalkyl, aryl, heterocycle,

5 unsubstituted or substituted with one or two:

a) C<sub>1-4</sub> alkoxy,

b) halogen, or

c) substituted or unsubstituted aryl or substituted or unsubstituted heterocycle;

10

R<sup>8</sup> is independently selected from hydrogen, C<sub>1-6</sub> alkyl, 2,2,2-trifluoroethyl, benzyl and aryl;

R<sup>9</sup> is independently selected from C<sub>1-6</sub> alkyl and aryl;

15

J, K and L are independently selected from: S, N, O, NH and CH;

X is a bond, -S(O)<sub>m</sub>-, O or -C(=O)-;

20 m is 0, 1 or 2;

p is 0, 1, 2, 3 or 4;


q is 0, 1 or 2, provided that q is not 0 or 1 if X is O; and

u is independently 0 or 1;

25 or an optical isomer or pharmaceutically acceptable salt thereof.

9. The compound according to Claim 7 which inhibits farnesyl-protein transferase of the formula G:






wherein:

5  $R^{1a}$  and  $R^{1c}$  are independently selected from: hydrogen,  $R^8O-$ ,  $-N(R^8)_2$ , F, C3-C10 cycloalkyl or C1-C6 alkyl;

$R^{1b}$  is independently selected from:

- 10 a) hydrogen,  
 b) aryl, heterocycle or C3-C10 cycloalkyl,  
 c) C1-C6 alkyl unsubstituted or substituted by aryl, heterocycle, C3-C10 cycloalkyl, C2-C6 alkenyl,  $R^8O-$ , or  $-N(R^8)_2$ ;

15  $R^{2a}$  and  $R^{2b'}$  are independently selected from selected from: H; C1-C6 alkyl,



$R^{3a}$  and  $R^{3b}$  are independently selected from:

- 20 a) hydrogen,  
 b) unsubstituted or substituted aryl, unsubstituted or substituted heterocycle, unsubstituted or substituted C3-C10 cycloalkyl, unsubstituted or substituted C2-C6 alkenyl, C2-C6 alkynyl, halogen, C1-C6 perfluoroalkyl,  $R^9O-$ ,  $R^9S(O)_m-$ ,  $R^8C(O)NR^8-$ ,  $(R^8)_2NC(O)-$ ,  $R^9C(O)O-$ ,

- $R^8_2N-C(NR^8)-$ , CN, NO<sub>2</sub>,  $R^8C(O)-$ , N<sub>3</sub>,  $-N(R^8)_2$ , or  $R^9OC(O)NR^8-$ ,  
 c) unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl,  
 d) substituted C<sub>1</sub>-C<sub>6</sub> alkyl wherein the substituent on the  
 5 substituted C<sub>1</sub>-C<sub>6</sub> alkyl is selected from unsubstituted or substituted aryl, unsubstituted or substituted heterocyclic, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl,  $R^9O-$ ,  $R^9S(O)_m-$ ,  $R^8C(O)NR^8-$ ,  $(R^8)_2NC(O)-$ ,  $R^8_2N-C(NR^8)-$ , CN,  $R^8C(O)-$ , N<sub>3</sub>,  $-N(R^8)_2$ , and  $R^9OC(O)-NR^8-$ ;  
 10  $R^{5a}$  and  $R^{5b}$  are independently hydrogen, ethyl, cyclopropyl or methyl;  
 $R^6$  and  $R^7$  are independently selected from:  
 H; C<sub>1</sub>-4 alkyl, C<sub>3</sub>-6 cycloalkyl, aryl, heterocycle,  
 15 unsubstituted or substituted with one or two:  
 a) C<sub>1</sub>-4 alkoxy,  
 b) halogen, or  
 c) substituted or unsubstituted aryl or substituted or unsubstituted heterocycle;  
 20  $R^8$  is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, 2,2,2-trifluoroethyl, benzyl and aryl;  
 $R^9$  is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;  
 25 J, K and L are independently selected from: S, N, O, NH and CH;  
 $A^1$  is selected from: a bond,  $-C(O)-$ , O,  $-N(R^8)-$ , or  $-S(O)_m$ ;  
 30 X is a bond,  $-S(O)_m-$ , O or  $-C(=O)-$ ;  
 m is 0, 1 or 2;  
 n is 0, 1 or 2; provided that n is not 0 if  $A^1$  is a bond, O,

-N(R<sup>8</sup>)-, or S(O)<sub>m</sub>;

p is 1, 2 or 3;

q is 0, 1 or 2, provided that q is not 0 or 1 if X is O; and

u is independently 0 or 1;


5

or an optical isomer or pharmaceutically acceptable salt thereof.

10. A compound which inhibits farnesyl-protein transferase which is:


10

5-(1-(4-Cyanobenzyl)-5-imidazolylmethyl)-4,5,6,7-tetrahydrothieno[3,2-c]pyridine




15

2-Bromo-5-(1-(4-cyanobenzyl)-5-imidazolylmethyl)-4,5,6,7-tetrahydrothieno[3,2-c]pyridine




20

2-Phenyl-5-(1-(4-cyanobenzyl)-5-imidazolylmethyl)-4,5,6,7-tetrahydrothieno[3,2-c]pyridine




25

5-[5-(5-Azabenzimidazolyl)methyl]-1-(4-cyanobenzyl)imidazole



5 5-[4-(4-Azabenzimidazolyl)methyl]-1-(4-cyanobenzyl)imidazole



or a pharmaceutically acceptable salt thereof.

11. A pharmaceutical composition comprising a  
10 pharmaceutical carrier, and dispersed therein, a therapeutically effective  
amount of a compound of Claim 1.

12. A pharmaceutical composition comprising a  
15 pharmaceutical carrier, and dispersed therein, a therapeutically effective  
amount of a compound of Claim 2.

13. A pharmaceutical composition comprising a  
20 pharmaceutical carrier, and dispersed therein, a therapeutically effective  
amount of a compound of Claim 4.

14. A pharmaceutical composition comprising a  
pharmaceutical carrier, and dispersed therein, a therapeutically effective  
amount of a compound of Claim 5.

15. A pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of Claim 10.

5                   16. A method for inhibiting farnesyl-protein transferase which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 11.

10                   17. A method for inhibiting farnesyl-protein transferase which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 12.

15                   18. A method for inhibiting farnesyl-protein transferase which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 13.

20                   19. A method for inhibiting farnesyl-protein transferase which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 14.

25                   20. A method for inhibiting farnesyl-protein transferase which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 15.

30                   21. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 11.

22. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 12.

5 23. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 13.

10 24. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 14.

15 25. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 15.

20 26. A method for treating neurofibromin benign proliferative disorder which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 12.

25 27. A method for treating blindness related to retinal vascularization which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 12.

28. A method for treating infections from hepatitis delta and related viruses which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 12.

30 29. A method for preventing restenosis which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 12.

30. A method for treating polycystic kidney disease which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 12.

5           31. A pharmaceutical composition made by combining the compound of Claim 1 and a pharmaceutically acceptable carrier.

32. A pharmaceutical composition made by combining the compound of Claim 2 and a pharmaceutically acceptable carrier.

10

33. A process for making a pharmaceutical composition comprising combining a compound of Claim 1 and a pharmaceutically acceptable carrier.

15

34. A process for making a pharmaceutical composition comprising combining a compound of Claim 2 and a pharmaceutically acceptable carrier.

## SEQUENCE LISTING

<110> Halczenko, Wasył  
Stump, Craig A.  
Merck & Co., Inc.

<120> INHIBITORS OF FARNESYL-PROTEIN  
TRANSFERASE

<130> 19849Y2

<150> 08/984,732

<151> 1997-12-04

<160> 26

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 4

<212> PRT

<213> Homosapien

<400> 1

Cys Val Ile Met

1

<210> 2

<211> 4

<212> PRT

<213> Homosapien

<400> 2

Cys Val Leu Leu

1

<210> 3

<211> 4

<212> PRT

<213> Homosapien

<400> 3

Cys Val Val Met

1

<210> 4

<211> 4

<212> PRT

<213> Homosapien

<400> 4

Cys Ile Ile Met

1

<210> 5



<211> 4  
<212> PRT  
<213> Homosapien

<400> 5  
Cys Leu Leu Leu  
1

<210> 6  
<211> 4  
<212> PRT  
<213> Homosapien

<400> 6  
Cys Gln Leu Leu  
1

<210> 7  
<211> 4  
<212> PRT  
<213> Homosapien

<400> 7  
Cys Ser Ile Met  
1

<210> 8  
<211> 4  
<212> PRT  
<213> Homosapien

<400> 8  
Cys Ala Ile Met  
1

<210> 9  
<211> 4  
<212> PRT  
<213> Homosapien

<400> 9  
Cys Lys Val Leu  
1

<210> 10  
<211> 4  
<212> PRT  
<213> Homosapien

<400> 10  
Cys Leu Ile Met  
1

<210> 11  
<211> 4  
<212> PRT

<213> Homosapien  
 <400> 11  
 Cys Val Leu Ser  
 1  
 <210> 12  
 <211> 4  
 <212> PRT  
 <213> Homosapien  
 <400> 12  
 Cys Ala Ile Leu  
 1  
 <210> 13  
 <211> 15  
 <212> PRT  
 <213> Homosapien  
 <400> 13  
 Gly Lys Lys Lys Lys Lys Lys Ser Lys Thr Lys Cys Val Ile Met  
 1 5 10 15  
 <210> 14  
 <211> 86  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> completely synthesized  
 <400> 14  
 tctcctcgag gccaccatgg ggagtagcaa gagcaagcct aaggacccca gccagcgccg 60  
 gatgacagaa tacaagcttg tgggtgg 86  
 <210> 15  
 <211> 33  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> completely synthesized  
 <400> 15 33  
 cacatctaga tcaggacagc acagacttgc agc  
 <210> 16  
 <211> 41  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> completely synthesized  
 <400> 16

tctcctogag gccaccatga cagaatacaa gcttgtggtg g 41

<210> 17  
<211> 38  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> completely synthesized

<400> 17  
cactctagac tgggtgcaga gcagcacaca cttgcagc 38

<210> 18  
<211> 38  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> completely synthesized

<400> 18  
gagagaattc gccaccatga cggaatataa gctggtgg 38

<210> 19  
<211> 33  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> completely synthesized

<400> 19  
gagagtcgac gcgtcaggag agcacacact tgc 33

<210> 20  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> completely synthesized

<400> 20  
ccgccggcct ggaggagtac ag 22

<210> 21  
<211> 38  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> completely synthesized

<400> 21  
gagagaattc gccaccatga ctgagtacaa actggtgg 38

<210> 22  
<211> 32  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> completely synthesized

<400> 22  
gagagtcgac ttgttacatc accacacatg gc

32

<210> 23  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> completely synthesized

<400> 23  
gttgagcag ttggtgttg g

21

<210> 24  
<211> 38  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> completely synthesized

<400> 24  
gagaggtacc gccacatga ctgaatataa acttggtg

38

<210> 25  
<211> 36  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> completely synthesized

<400> 25  
ctctgtcgac gtatttacat aattacacac ttgtgc

36

<210> 26  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> completely synthesized

<400> 26  
gtagttggag ctggtggcgt aggc

24

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/25324

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) :A61K 31/44; C07D 471/04

US CL :546/114, 118; 514/301, 303

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 546/114, 118; 514/301, 303

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Chemical Abstracts on STN

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE 42 30 464 A1 (MEDERSKI et al) 17 March 1994, page 2, lines 1-48.	1-34

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

01 FEBRUARY 1999

Date of mailing of the international search report

11 FEB 1999

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ANN RAZGUNAS

Telephone No. (703) 308-1235

**THIS PAGE BLANK (USPTO)**